

Neutrino mass and mixing in the 3-3-1 model and S_3 flavor symmetry with minimal Higgs content

V. V. Vien*

*Department of Physics, Tay Nguyen University,
567 Le Duan, Buon Ma Thuot, DakLak, Vietnam*

H. N. Long†

Institute of Physics, VAST, 10 Dao Tan, Ba Dinh, Hanoi, Vietnam

(Dated: July 11, 2014)

A new S_3 flavor model based on $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$ gauge symmetry responsible for fermion masses and mixings different from our previous work [12, 13] is constructed. The new feature is a two - dimensional representation of a Higgs anti-sextet under S_3 which responsible for neutrino masses and mixings. The neutrinos acquire small masses from only an anti-sextet of $SU(3)$ which is in a doublet under S_3 . If the difference of components of the anti-sextet is regarded as a small perturbation, S_3 is equivalently broken into identity, the corresponding neutrino mass mixing matrix acquires the most general form and the model can fit the latest data on neutrino oscillation. This way of the symmetry breaking helps us to reduce a content in the Higgs sector, only one an anti-sextet instead of two as in our previous work [13]. Our results show that the neutrino masses are naturally small and a small deviation from the tri-bimaximal neutrino mixing form can be realized. The Higgs potential of the model as well as the minimization conditions and gauge boson masses and mixings is also considered.

PACS numbers: 14.60.Pq, 14.60.St, 12.60.Fr, 11.30.Er

I. INTRODUCTION

The experiments of neutrino oscillations have indicated that the neutrinos have small masses and mixings [1–4], and therefore the standard model of fundamental particles and interactions must be extended. Among this direction, there have been various models proposed, such as [5, 6] and others. An alternative is to extend the electroweak symmetry $SU(2)_L \otimes U(1)_Y$ into $SU(3)_L \otimes U(1)_X$, in which to complete the fundamental representations of $SU(3)_L$ with the standard-model doublets

*Electronic address: wvienk16@gmail.com

†Electronic address: hnlong@iop.vast.ac.vn

so as to obtain the neutral fermions. This proposal has nice features and has been extensively studied over the last two decades, is called 3-3-1 models [7, 9, 10], in which the number of fermion families has been proven to be three [7, 8].

The parameters of neutrino oscillations such as the squared mass differences and mixing angles are now very constrained. The data in Ref.[4] imply that

$$\begin{aligned}\sin^2(2\theta_{12}) &= 0.857 \pm 0.024 \quad (t_{12} \simeq 0.6717), \\ \sin^2(2\theta_{13}) &= 0.098 \pm 0.013 \quad (s_{13} \simeq 0.1585), \\ \sin^2(2\theta_{23}) &> 0.95, \\ \Delta m_{21}^2 &= (7.50 \pm 0.20) \times 10^{-5} \text{eV}^2, \quad \Delta m_{32}^2 = (2.32^{+0.12}_{-0.08}) \times 10^{-3} \text{eV}^2.\end{aligned}\tag{1}$$

These large neutrino mixing angles are completely different from the quark mixing ones defined by the CKM matrix. Therefore, it is very important to find a natural model that leads to these mixing patterns of quarks and leptons with good accuracy. Small non-Abelian discrete symmetries are considered to be the most attractive choice for the flavor sector [15–17]. The simplest explanation for these conclusions is probably due to an S_3 flavor symmetry which is the smallest non-Abelian discrete group [13, 20]. In fact, there is an approximately maximal mixing of two flavors μ and τ as given above which can be connected by a $\underline{2}$ irreducible representation of S_3 . Besides the $\underline{2}$, the group S_3 can provide two inequivalent singlet representations $\underline{1}$ and $\underline{1}'$ which play a crucial role in reproducing consistent fermion masses and mixings [13]. The S_3 models have been studied extensively over the last decade [17]. In [13] we have proposed two 3-3-1 models, with either neutral fermions or right-handed neutrinos, based on S_3 flavor symmetry, in which there is a large number of Higgs triplets was required. In this paper, we propose a new S_3 flavor symmetry in the 3-3-1 model with neutral fermions, in which the number of Higgs triplets required is less and the Higgs potential of the model is therefore much simpler than the previous ones.

The motivation for extending the above application to the 3-3-1 models with the neutral fermions N_R is mentioned in [11–13]. In this paper, we investigate simpler choices for Higgs multiples of S_3 in which the unique anti-sextet responsible for neutrino mass and mixing lying in $\underline{2}$ under S_3 and the difference between two VEV components of anti-sextet plays the role of perturbation. It is also noted that the numbers of fermion families in the 3-3-1 models have an origin from the anomaly-free gauge symmetry naturally meet our criteria on the dimensions of flavor group representations as such S_3 , unlike the others in the literature, mostly imposed by hand [15–17].

The rest of this work is as follows. In Sec. II we present the necessary elements of the 3-3-1 model with neutral fermions N_R under the S_3 symmetry and introduce the necessary Higgs

fields responsible for the charged-lepton and quark masses. Section III is devoted to the neutrino mass and mixing. In Sec. IV we consider the Higgs potential and minimization conditions. We summarize our results and make conclusions in the section VI.

II. THE MODEL

The fermion content of the model is similar to that in [13]: the fermions in the model transform under respective $[SU(3)_L, U(1)_X, U(1)_{\mathcal{L}}, \underline{S}_3]$ symmetries as

$$\begin{aligned}
\psi_{1L} &= (\nu_{1L}, l_{1L}, N_{1R}^c)^T \sim [3, -1/3, 2/3, \underline{1}], \quad l_{1R} \sim [1, -1, 1, \underline{1}], \\
\psi_{\alpha L} &= (\nu_{\alpha L}, l_{\alpha L}, N_{\alpha R}^c)^T \sim [3, -1/3, 2/3, \underline{2}], \quad l_{\alpha R} \sim [1, -1, 1, \underline{2}], \\
Q_{1L} &= (u_{1L}, d_{1L}, U_L)^T \sim [3, 1/3, -1/3, \underline{1}], \\
u_{1R} &\sim [1, 2/3, 0, \underline{1}], \quad d_{1R} \sim [1, -1/3, 0, \underline{1}], \quad U_R \sim [1, 2/3, -1, \underline{1}], \\
Q_{\alpha L} &= (d_{\alpha L}, -u_{\alpha L}, D_{\alpha L})^T \sim [3^*, 0, 1/3, \underline{2}], \\
u_{\alpha R} &\sim [1, 2/3, 0, \underline{2}], \quad d_{\alpha R} \sim [1, -1/3, 0, \underline{2}], \quad D_{\alpha R} \sim [1, -1/3, 1, \underline{2}].
\end{aligned} \tag{2}$$

where $\alpha = 2, 3$ is a family index of the last two lepton and quark families, which are defined as the components of the $\underline{2}$ representations. We note that the $\underline{2}$ for quarks satisfies the requirement of anomaly cancelation, where the last two left-quark families are in 3^* and the first one as well as the leptons are in 3 . All the \mathcal{L} charges of the model multiplets are listed in the square brackets. In what follows, we consider possibilities for generating the fermion masses. The scalar multiplets needed for this purpose are to be introduced accordingly.

To generate masses for the charged leptons, we introduce two $SU(3)_L$ scalar triplets ϕ and ϕ' respectively lying in $\underline{1}$ and $\underline{1}'$ under S_3 , with the VEVs $\langle \phi \rangle = (0 \ v \ 0)^T$ and $\langle \phi' \rangle = (0 \ v' \ 0)^T$ [13]. From the invariant Yukawa interactions for the charged leptons, we obtain $m_e = h_1 v$, $m_\mu = hv - h'v'$, $m_\tau = hv + h'v'$, and the left and right-handed charged leptons mixing matrices are diagonal, $U_{lL} = U_{lR} = 1$. The charged leptons $l_{1,2,3}$ therefore by themselves are the physical mass eigenstates and the lepton mixing matrix depends on only that of the neutrinos, which is studied in the next section.

In similarity to the charged lepton sector, to generate the quark masses, we additionally introduce the three scalar Higgs triplets χ , η , η' respectively lying in $\underline{1}$, $\underline{1}$ and $\underline{1}'$ under S_3 . Quark masses can be derived from the invariant Yukawa interactions for quarks, assuming that the VEVs of η , η' and χ are u , u' and w , where $u = \langle \eta_1^0 \rangle$, $u' = \langle \eta_1'^0 \rangle$, and $w = \langle \chi_3^0 \rangle$ and the other VEVs $\langle \eta_3^0 \rangle$,

$\langle \eta_3^0 \rangle$, and $\langle \chi_1^0 \rangle$ vanish due to the lepton parity conservation. The exotic quarks therefore acquire masses $m_U = f_1 w$ and $m_{D_{1,2}} = fw$. The masses of ordinary up-quarks and down-quarks are

$$\begin{aligned} m_u &= h_1^u u, & m_c &= h^u v + h'^u v', & m_t &= h^u v - h'^u v', \\ m_d &= h_1^d v, & m_s &= h^d u + h'^d u', & m_b &= h^d u - h'^d u'. \end{aligned}$$

The unitary matrices which couple the left-handed quarks u_L and d_L to those in the mass bases are unit ones. The CKM quark mixing matrix at the tree level is then $U_{\text{CKM}} = U_{dL}^\dagger U_{uL} = 1$. The lepton parity breaking due to the odd VEVs $\langle \eta_3^0 \rangle$, $\langle \eta_3'^0 \rangle$, $\langle \chi_1^0 \rangle$, or a violation of \mathcal{L} and/or S_3 symmetry in terms of Yukawa interactions would disturb the tree-level matrix, resulting in a mixing between the SM and exotic quarks and/or possibly providing the desirable quark mixing pattern $\bar{Q}_{1L}\chi_{u1R}$, $\bar{Q}_L\chi^*d_R$, $\bar{Q}_{1L}\chi_{uR}$, with a mixing between SM and exotic quarks. To obtain a realistic pattern of the SM quarks mixing, we should add radiative correction or use the effective six-dimensional operators (see Ref. [18] for details). However, we leave this problem for the future work. A detailed study on charged lepton and quark masses can be found in Ref. [13]. In this paper, we consider a new representation for the anti-sextet responsible for neutrino masses and mixings that are different from those in Ref. [13].

III. NEUTRINO MASSES AND MIXING

The neutrino masses arise from the couplings of $\bar{\psi}_{\alpha L}^c \psi_{\alpha L}$, $\bar{\psi}_{1L}^c \psi_{1L}$ and $\bar{\psi}_{1L}^c \psi_{\alpha L}$ to scalars, where $\bar{\psi}_{\alpha L}^c \psi_{\alpha L}$ transforms as $3^* \oplus 6$ under $\text{SU}(3)_L$ and as $\underline{1} \oplus \underline{1}' \oplus \underline{2}$ under S_3 ; $\bar{\psi}_{1L}^c \psi_{1L}$ transforms as $3^* \oplus 6$ under $\text{SU}(3)_L$ and as $\underline{1}$ under S_3 , and $\bar{\psi}_{1L}^c \psi_{\alpha L}$ transforms as $3^* \oplus 6$ under $\text{SU}(3)_L$ and as $\underline{2}$ under S_3 . For the known scalar triplets $(\phi, \phi', \chi, \eta, \eta')$, the available interactions are only $(\bar{\psi}_{\alpha L}^c \psi_{\alpha L})\phi$ and $(\bar{\psi}_{\alpha L}^c \psi_{\alpha L})\phi'$, but are explicitly suppressed because of the \mathcal{L} -symmetry. We therefore propose a new $\text{SU}(3)_L$ antisextet coupling to $\bar{\psi}_L^c \psi_L$ responsible for the neutrino masses lying in either $\underline{1}$, $\underline{1}'$, or $\underline{2}$ under S_3 . To obtain a realistic neutrino spectrum with minimal Higgs content, we introduce the Higgs anti-sextet

$$s_i = \begin{pmatrix} s_{11}^0 & s_{12}^+ & s_{13}^0 \\ s_{12}^+ & s_{22}^{++} & s_{23}^+ \\ s_{13}^0 & s_{23}^+ & s_{33}^0 \end{pmatrix}_i \sim [6^*, 2/3, -4/3, \underline{2}], \quad (i = 1, 2)$$

where the numbered subscripts on the component scalars are the $SU(3)_L$ indices, whereas $i = 1, 2$ is that of S_3 . The VEV of s is set as $(\langle s_1 \rangle, \langle s_2 \rangle)$ under S_3 , with

$$\langle s_i \rangle = \begin{pmatrix} \lambda_i & 0 & v_i \\ 0 & 0 & 0 \\ v_i & 0 & \Lambda_i \end{pmatrix}, \quad (i = 1, 2). \quad (3)$$

Following the potential minimization conditions, we have several VEV alignments. The first one is that $\langle s_1 \rangle = \langle s_2 \rangle$; then S_3 is broken into Z_2 consisting of the identity element and one transposition (out of the three) of S_3 . The second one is that $\langle s_1 \rangle \neq 0 = \langle s_2 \rangle$ or $\langle s_1 \rangle = 0 \neq \langle s_2 \rangle$; then S_3 is broken into Z_3 as in the case of the charged lepton sector. The third one is that $\langle s_1 \rangle \neq \langle s_2 \rangle$; then S_3 is broken into the identity. In our previous work [13], we have argued that both breakings $S_3 \rightarrow Z_2$ and $S_3 \rightarrow Z_3$ must take place, and hence, to obtain a realistic neutrino spectrum, we additionally introduced a triplet (ρ) and an anti-sextet (s) that lie in $\underline{1}'$ and $\underline{2}$ under S_3 . With these alignments, the number of Higgs multiplets required is eight. In this work, we propose that both the first and the third direction take place. The Yukawa interactions are

$$\begin{aligned} -\mathcal{L}_\nu &= \frac{x}{2}(\bar{\psi}_{1L}^c s)\underline{2}\psi_{\alpha L} + \frac{y}{2}(\bar{\psi}_{\alpha L}^c s)\underline{2}\psi_{\alpha L} + h.c. \\ &= \frac{x}{2}\bar{\psi}_{1L}^c(\psi_{2L}s_2 + \psi_{3L}s_1) + \frac{y}{2}(\bar{\psi}_{2L}^c\psi_{2L}s_1 + \bar{\psi}_{3L}^c\psi_{3L}s_2) + h.c., \end{aligned} \quad (4)$$

where the Yukawa coupling x is that of lepton flavor changing interactions. The mass Lagrangian for the neutrinos is given by

$$\begin{aligned} -\mathcal{L}_\nu^{\text{mass}} &= \frac{1}{2}x(\lambda_2\bar{\nu}_{1L}^c\nu_{2L} + v_2\bar{\nu}_{1L}^cN_{2R}^c + v_2\bar{N}_{1R}\nu_{2L} + \Lambda_2\bar{N}_{1R}N_{2R}^c) \\ &+ \frac{1}{2}x(\lambda_1\bar{\nu}_{1L}^c\nu_{3L} + v_1\bar{\nu}_{1L}^cN_{3R}^c + v_1\bar{N}_{1R}\nu_{3L} + \Lambda_1\bar{N}_{1R}N_{3R}^c) \\ &+ \frac{1}{2}y(\lambda_1\bar{\nu}_{2L}^c\nu_{2L} + v_1\bar{\nu}_{2L}^cN_{2R}^c + v_1\bar{N}_{2R}\nu_{2L} + \Lambda_1\bar{N}_{2R}N_{2R}^c) \\ &+ \frac{1}{2}y(\lambda_2\bar{\nu}_{3L}^c\nu_{3L} + v_2\bar{\nu}_{3L}^cN_{3R}^c + v_2\bar{N}_{3R}\nu_{3L} + \Lambda_2\bar{N}_{3R}N_{3R}^c) + h.c. \end{aligned} \quad (5)$$

and also by

$$-\mathcal{L}_\nu^{\text{mass}} = \frac{1}{2}\bar{\chi}_L^c M_\nu \chi_L + h.c., \quad \chi_L \equiv \begin{pmatrix} \nu_L \\ N_R^c \end{pmatrix}, \quad M_\nu \equiv \begin{pmatrix} M_L & M_D^T \\ M_D & M_R \end{pmatrix}, \quad (6)$$

where $\nu = (\nu_1, \nu_2, \nu_3)^T$ and $N = (N_1, N_2, N_3)^T$. The mass matrices are then obtained by

$$M_{L,R,D} = \begin{pmatrix} 0 & a_{L,R,D} & b_{L,R,D} \\ a_{L,R,D} & c_{L,R,D} & 0 \\ b_{L,R,D} & 0 & d_{L,R,D} \end{pmatrix}, \quad (7)$$

with

$$\begin{aligned}
a_L &= \frac{x}{2}\lambda_s \equiv \frac{x}{2}\lambda_2, \quad a_D = \frac{x}{2}v_s \equiv \frac{x}{2}v_2, \quad a_R = \frac{x}{2}\Lambda_s \equiv \frac{x}{2}\Lambda_2, \\
b_L &= \frac{x}{2}\lambda_1, \quad b_D = \frac{x}{2}v_1, \quad b_R = \frac{x}{2}\Lambda_1, \\
c_L &= y\lambda_1, \quad c_D = yv_1, \quad c_R = y\Lambda_1, \\
d_L &= y\lambda_s \equiv y\lambda_2, \quad d_D = yv_s \equiv yv_2, \quad d_R = y\Lambda_s \equiv y\Lambda_2.
\end{aligned} \tag{8}$$

In general, three active-neutrinos therefore gain masses via a combination of type I and type II seesaw mechanisms, derived from (6) and (7) as

$$M_{\text{eff}} = M_L - M_D^T M_R^{-1} M_D = \begin{pmatrix} A & B_1 & B_2 \\ B_1 & C_1 & D \\ B_2 & D & C_2 \end{pmatrix}, \tag{9}$$

where

$$\begin{aligned}
A &= -\frac{(a_R b_D - a_D b_R)^2}{b_R^2 c_R + a_R^2 d_R}, \\
B_1 &= \frac{b_R [a_R b_D c_D + a_L b_R c_R - a_D (b_R c_D + b_D c_R)] + a_R (a_L a_R - a_D^2) d_R}{b_R^2 c_R + a_R^2 d_R}, \\
B_2 &= \frac{-b_D^2 b_R c_R + b_L b_R^2 c_R + a_D a_R b_R d_D + a_R^2 b_L d_R - a_R b_D (a_R d_D + a_D d_R)}{b_R^2 c_R + a_R^2 d_R}, \\
C_1 &= \frac{b_R^2 (c_L c_R - c_D^2) + (a_R^2 c_L + a_D^2 c_R - 2a_D a_R c_D) d_R}{b_R^2 c_R + a_R^2 d_R}, \\
C_2 &= \frac{-2b_D b_R c_R d_D + b_R^2 c_R d_L + b_D^2 c_R d_R + a_R^2 (d_L d_R - d_D^2)}{b_R^2 c_R + a_R^2 d_R}, \\
D &= \frac{(a_R c_D - a_D c_R)(b_R d_D - b_D d_R)}{b_R^2 c_R + a_R^2 d_R}.
\end{aligned} \tag{10}$$

The neutrino mass matrix in (9) is similar to the one in Ref.[13] but the broken symmetry directions are difference. Indeed, in this model there are two broken symmetry directions as follows.

- If S_3 is broken to Z_2 (the subgroup Z_2 is unbroken), then we have $A = D = 0$, $B_1 = B_2$ and $C_1 = C_2$.
- If $S_3 \rightarrow \{\text{Identity}\}$ (or, equivalently, $Z_2 \rightarrow \{\text{Identity}\}$), then we have $A \neq 0$, $D \neq 0$, $B_1 \neq B_2$ and $C_1 \neq C_2$, but A and D are close to zero, and B_1, B_2, C_1, C_2 kept close to each other in pairs. In this case, the disparity between $\langle s_1 \rangle$ and $\langle s_2 \rangle$ is very small and can be regarded as a small perturbation.

We next divide our considerations into two cases to fit the data: the first case is where only S_3 is broken to Z_2 , and the second case is a combination of the both $S_3 \rightarrow Z_2$ and $Z_2 \rightarrow \{\text{Identity}\}$.

A. Experimental constraints under $S_3 \rightarrow Z_2$

In the case $S_3 \rightarrow Z_2$, $\lambda_1 = \lambda_2 \equiv \lambda_s$, $v_1 = v_2 \equiv v_s$, $\Lambda_1 = \Lambda_2 \equiv \Lambda_s$, we have $A = D = 0$, $B_1 = B_2 \equiv B$, $C_1 = C_2 \equiv C$, and M_{eff} in (9) reduces to

$$M_{\text{eff}} = \begin{pmatrix} 0 & B & B \\ B & C & 0 \\ B & 0 & C \end{pmatrix}, \quad (11)$$

with

$$B = \left(\lambda_s - \frac{v_s^2}{\Lambda_s} \right) \frac{x}{2}, \quad C = \left(\lambda_s - \frac{v_s^2}{\Lambda_s} \right) y. \quad (12)$$

We can diagonalize the matrix M_{eff} in (11) as

$$U^T M_{\text{eff}} U = \text{diag}(m_1, m_2, m_3),$$

where

$$\begin{aligned} m_1 &= \frac{1}{2} \left(C - \sqrt{C^2 + 8B^2} \right) = \left(\lambda_s - \frac{v_s^2}{\Lambda_s} \right) \frac{y + \sqrt{y^2 + 2x^2}}{2}, \\ m_2 &= \frac{1}{2} \left(C + \sqrt{C^2 + 8B^2} \right) = \left(\lambda_s - \frac{v_s^2}{\Lambda_s} \right) \frac{y - \sqrt{y^2 + 2x^2}}{2}, \\ m_3 &= C = \left(\lambda_s - \frac{v_s^2}{\Lambda_s} \right) y, \end{aligned} \quad (13)$$

and the neutrino mixing matrix takes the form

$$U_0 = \begin{pmatrix} \frac{|K|}{\sqrt{|K|^2+2}} & -\frac{\sqrt{2}}{\sqrt{|K|^2+2}} & 0 \\ \frac{1}{\sqrt{|K|^2+2}} & \frac{1}{\sqrt{2}} \frac{|K|}{\sqrt{|K|^2+2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{|K|^2+2}} & \frac{1}{\sqrt{2}} \frac{|K|}{\sqrt{|K|^2+2}} & \frac{1}{\sqrt{2}} \end{pmatrix}, \quad K = -\frac{C + \sqrt{C^2 + 8B^2}}{2B} \quad (14)$$

We note that $m_1 m_2 = -2B^2$. This matrix can be parameterized by three Euler's angles, which implies

$$\theta_{13} = 0, \quad \theta_{23} = \pi/4, \quad \tan \theta_{12} = \frac{\sqrt{2}}{|K|}. \quad (15)$$

This case coincides with the data because $\sin^2(2\theta_{13}) < 0.15$ and $\sin^2(2\theta_{23}) > 0.92$ [3]. For the remaining constraints, taking the central values from the data [3],

$$\begin{aligned} \sin^2(2\theta_{12}) &\simeq 0.87, \quad (s_{12}^2 = 0.32), \\ \Delta m_{21}^2 &= 7.59 \times 10^{-5} \text{ eV}^2, \quad \Delta m_{32}^2 = 2.43 \times 10^{-3} \text{ eV}^2, \end{aligned}$$

we have a solution

$$m_1 = 0.0280284 \text{ eV}, \quad m_2 = 0.0293347 \text{ eV}, \quad m_3 = 0.0573631 \text{ eV}, \quad (16)$$

and $B = -0.0202757i \text{ eV}$, $C = 0.0573631 \text{ eV}$, $K = 1.44667$, $|x/y| = 0.707087$. It follows that $\tan \theta_{12} = 0.977565$, $(\theta_{12} \simeq 44.35^\circ)$, and the neutrino mixing matrix form is very close to that of the bi-maximal mixing pattern mentioned in Ref. [21]:

$$U = \begin{pmatrix} 0.715083 & -0.69904 & 0 \\ 0.494296 & 0.50564 & -\frac{1}{\sqrt{2}} \\ 0.494296 & 0.50564 & \frac{1}{\sqrt{2}} \end{pmatrix} \cong \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{\sqrt{2}} \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (17)$$

Now, it is natural to choose $\lambda_s, v_s^2/\Lambda_s$ in eV order, and suppose that $\lambda_s > v_s^2/\Lambda_s$. We assume that $\lambda_s - v_s^2/\Lambda_s = 0.1$, we have $x = -0.573631$ and $y = 0.399403i$.

It was assumed in recent analyses that $\theta_{13} \neq 0$, but is small, as in Ref.[4]. If this is correct, then that case will fail. But the direction of the breakings $S_3 \rightarrow \{\text{Identity}\}$ can improve this.

B. Experimental constraints under $S_3 \rightarrow \{\text{Identity}\}$

If both $S_3 \rightarrow Z_2$ and $Z_2 \rightarrow \{\text{Identity}\}$ directions are realized, then $\lambda_1 \neq \lambda_2 \equiv \lambda_s$, $v_1 \neq v_2 \equiv v_s$ and $\Lambda_1 \neq \Lambda_2 \equiv \Lambda_s$ and, consequently $A \approx 0$, $D_1 \approx 0$, $B_1 \approx B_2$ and $C_1 \approx C_2$, and the general neutrino mass matrix in (9) can be rewritten in the form

$$M_{\text{eff}} = \begin{pmatrix} 0 & B & B \\ B & C & 0 \\ B & 0 & C \end{pmatrix} + \begin{pmatrix} r_1 & p_1 & p_2 \\ p_1 & q_1 & r_2 \\ p_2 & r_2 & q_2 \end{pmatrix}. \quad (18)$$

where B and C are given by (12), to match the case $S_2 \rightarrow Z_2$ as in (11). The last matrix in (18) is a deviation from the contribution due to the disparity of $\langle s_1 \rangle$ and $\langle s_2 \rangle$, namely

$$p_1 = B_1 - B, \quad B_2 - B = p_1, \quad C_1 - C = q_1, \quad C_2 - C = q_2, \quad r_1 = A, \quad r_2 = D. \quad (19)$$

With the A, D and $B_{1,2}, C_{1,2}$ defined in (10), it corresponds to $S_3 \rightarrow \{\text{Identity}\}$. Substituting (10) and (12) into (19) with the help of (8), we obtain

$$r_1 = -\frac{(\Lambda_s v_1 - \Lambda_1 v_s)^2 x^2}{\Lambda_1^3 + \Lambda_s^3} \frac{x^2}{4y} = -\frac{\Lambda_1^2 \Lambda_s^2}{\Lambda_1^3 + \Lambda_s^3} \left(\frac{v_1}{\Lambda_1} - \frac{v_s}{\Lambda_s} \right)^2 \frac{x^2}{4y}, \quad (20)$$

$$r_2 = -\frac{(\Lambda_s v_1 - \Lambda_1 v_s)^2 y}{\Lambda_1^3 + \Lambda_s^3} = -\frac{\Lambda_1^2 \Lambda_s^2}{\Lambda_1^3 + \Lambda_s^3} \left(\frac{v_1}{\Lambda_1} - \frac{v_s}{\Lambda_s} \right)^2 y, \quad (21)$$

$$p_1 = \frac{\Lambda_1(\Lambda_s v_1 - \Lambda_1 v_s)^2 x}{2\Lambda_s(\Lambda_1^3 + \Lambda_s^3)} = \frac{\Lambda_1}{\Lambda_s} \frac{\Lambda_1^2 \Lambda_s^2}{\Lambda_1^3 + \Lambda_s^3} \left(\frac{v_1}{\Lambda_1} - \frac{v_s}{\Lambda_s} \right)^2 \frac{x}{2}, \quad (22)$$

$$p_2 = \frac{(\lambda_1 - \lambda_s)\Lambda_s(\Lambda_1^3 + \Lambda_s^3) - \Lambda_1^2 \Lambda_s v_1^2 - 2\Lambda_s^3 v_1 v_s + (\Lambda_1^3 + \Lambda_1 \Lambda_s^2 + \Lambda_s^3) x}{\Lambda_s(\Lambda_1^3 + \Lambda_s^3) v_s^2} \frac{x}{2}, \quad (23)$$

$$\begin{aligned} q_1 &= \frac{[(\lambda_1 - \lambda_s)\Lambda_s(\Lambda_1^3 + \Lambda_s^3) - \Lambda_1^2 \Lambda_s v_1^2 - 2\Lambda_s^3 v_1 v_s + (\Lambda_1^3 + \Lambda_1 \Lambda_s^2 + \Lambda_s^3) v_s^2] y}{\Lambda_s(\Lambda_1^3 + \Lambda_s^3)} \\ &= (\lambda_1 - \lambda_s) y - \frac{(\Lambda_1^2 \Lambda_s v_1^2 + 2\Lambda_s^3 v_1 v_s) y}{\Lambda_s(\Lambda_1^3 + \Lambda_s^3)} + \frac{(\Lambda_1^3 + \Lambda_1 \Lambda_s^2 + \Lambda_s^3) v_s^2 y}{\Lambda_s(\Lambda_1^3 + \Lambda_s^3)} \\ &= (\lambda_1 - \lambda_s) y - \frac{(v_1^2 + 2\frac{\Lambda_s^2}{\Lambda_1^2} v_1 v_s) y}{\Lambda_1 + \frac{\Lambda_s^2}{\Lambda_1^2} \Lambda_s} + \frac{\left(\frac{\Lambda_1}{\Lambda_s} + \frac{\Lambda_s}{\Lambda_1} + \frac{\Lambda_s^2}{\Lambda_1^2} \right) v_s^2 y}{\Lambda_1 + \frac{\Lambda_s^2}{\Lambda_1^2} \Lambda_s}, \end{aligned} \quad (24)$$

$$q_2 = \frac{\Lambda_1(\Lambda_s v_1 - \Lambda_1 v_s)^2 y}{\Lambda_s(\Lambda_1^3 + \Lambda_s^3)} = \frac{\Lambda_1}{\Lambda_s} \frac{\Lambda_1^2 \Lambda_s^2}{\Lambda_1^3 + \Lambda_s^3} \left(\frac{v_1}{\Lambda_1} - \frac{v_s}{\Lambda_s} \right)^2 y. \quad (25)$$

Indeed, if $S_3 \rightarrow Z_2$, then the deviations p_i, q_i and r_i ($i = 1, 2$) vanish, and therefore the mass matrix M_{eff} in (9) reduces to its first term coinciding with (11). The first term in (18) provides a bi-maximal mixing pattern with $\theta_{13} = 0$ as shown in Sect.III A. The others, proportional to p_i, q_i, r_i due to contribution of the disparity of $\langle s_1 \rangle$ and $\langle s_2 \rangle$ take the role of perturbation for such a deviation of θ_{13} . Hence, in this work we consider the disparity of $\langle s_1 \rangle$ and $\langle s_2 \rangle$ contribution as a small perturbation and truncate the theory at the first order.

In Ref. [19] we considered the case of $S_4 \rightarrow K_4$ breaking corresponding to $S_3 \rightarrow \{\text{Identity}\}$ with $\lambda_1 \neq \lambda_s$ but $v_1 = v_s$ and $\Lambda_1 = \Lambda_s$. Then $r_1 = r_2 = p_1 = q_2 = 0$, $p_2 = \frac{x}{2y} q_1$, $q_1 = (\lambda_1 - \lambda_s) y \equiv \epsilon y$ with $\epsilon = \lambda_1 - \lambda_s$ being a small parameter that plays the role of a perturbation. In this paper, we consider the more general case, in which all elements of $\langle s_1 \rangle$ and $\langle s_2 \rangle$ are different from each other.

If $|\langle s_1 - s_2 \rangle| \ll \langle s_1 \rangle \sim \langle s_2 \rangle$ and $\frac{v_1}{\Lambda_1} \sim \frac{v_s}{\Lambda_s} \ll 1$, then we can evaluate $r_1, r_2, p_1, q_2 \ll 1$ which are of the second order in the perturbation and are therefore ignored. The remaining parameters p_2, q_1 are easily obtained as

$$p_2 = \alpha \frac{x}{2}, \quad q_1 = \alpha y, \quad (26)$$

where

$$\alpha = (\lambda_1 - \lambda_s) - \frac{(v_1^2 + 2\frac{\Lambda_s^2}{\Lambda_1^2} v_1 v_s)}{\Lambda_1 + \frac{\Lambda_s^2}{\Lambda_1^2} \Lambda_s} + \frac{\left(\frac{\Lambda_1}{\Lambda_s} + \frac{\Lambda_s}{\Lambda_1} + \frac{\Lambda_s^2}{\Lambda_1^2} \right) v_s^2}{\Lambda_1 + \frac{\Lambda_s^2}{\Lambda_1^2} \Lambda_s}. \quad (27)$$

The matrix M_{eff} in (18) thus reduces to

$$M_{\text{eff}} = \begin{pmatrix} 0 & B & B \\ B & C & 0 \\ B & 0 & C \end{pmatrix} + \alpha \begin{pmatrix} 0 & 0 & \frac{x}{2} \\ 0 & y & 0 \\ \frac{x}{2} & 0 & 0 \end{pmatrix} \equiv M_{\text{eff}}^0 + \alpha M^{(1)}. \quad (28)$$

Evaluating α shows that it is a small parameter, which can be regarded as a small perturbation. Within the perturbation theory up to the first order of α , the physical neutrino masses are obtained as

$$\begin{aligned} m'_1 &= \lambda_1 = m_1 + \alpha \left(\frac{Kx + y}{K^2 + 2} \right), \\ m'_2 &= \lambda_2 = m_2 + \frac{\alpha K(Ky - 2x)}{2(K^2 + 2)}, \quad m'_3 = \lambda_3 = m_3 + \alpha \frac{y}{2}, \end{aligned} \quad (29)$$

where $m_{1,2,3}$ are the mass values as in the case $S_3 \rightarrow Z_2$ given by (16). For the corresponding perturbed eigenstates, we set

$$U \longrightarrow U' = U + \Delta U,$$

where U is defined by (14), and

$$\Delta U = \begin{pmatrix} \Delta U_{11} & \Delta U_{12} & \Delta U_{13} \\ \Delta U_{21} & \Delta U_{22} & \Delta U_{23} \\ \Delta U_{31} & \Delta U_{32} & \Delta U_{33} \end{pmatrix}, \quad (30)$$

where

$$\begin{aligned} \Delta U_{11} &= -\alpha \frac{(K^2 - 2)x + 2Ky}{2(K^2 + 2)^{\frac{3}{2}}(m_1 - m_2)}, \\ \Delta U_{21} &= -\alpha \frac{(Kx - 2y)}{4\sqrt{K^2 + 2}(m_1 - m_3)} + \alpha \frac{K[(K^2 - 2)x + 2Ky]}{4(K^2 + 2)^{\frac{3}{2}}(m_1 - m_2)}, \\ \Delta U_{31} &= \alpha \frac{(Kx - 2y)}{4\sqrt{K^2 + 2}(m_1 - m_3)} + \alpha \frac{K[(K^2 - 2)x + 2Ky]}{4(K^2 + 2)^{\frac{3}{2}}(m_1 - m_2)}, \\ \Delta U_{12} &= -\alpha \frac{K[(K^2 - 2)x + 2Ky]}{2\sqrt{2}(K^2 + 2)^{3/2}(m_1 - m_2)}, \\ \Delta U_{22} &= \frac{\alpha}{2\sqrt{2}} \frac{Ky + x}{\sqrt{K^2 + 2}(m_2 - m_3)} - \frac{\alpha}{2\sqrt{2}} \frac{(K^2 - 2)x + 2Ky}{(K^2 + 2)^{\frac{3}{2}}(m_1 - m_2)}, \\ \Delta U_{32} &= -\frac{\alpha}{2\sqrt{2}} \frac{Ky + x}{\sqrt{K^2 + 2}(m_2 - m_3)} - \frac{\alpha}{2\sqrt{2}} \frac{(K^2 - 2)x + 2Ky}{(K^2 + 2)^{\frac{3}{2}}(m_1 - m_2)}, \\ \Delta U_{13} &= -\frac{\alpha}{2\sqrt{2}} \frac{K(Kx - 2y)}{(K^2 + 2)(m_1 - m_3)} - \frac{\alpha}{\sqrt{2}} \frac{Ky + x}{(K^2 + 2)(m_2 - m_3)}, \\ \Delta U_{23} &= \Delta U_{33} = -\frac{\alpha}{2\sqrt{2}} \frac{Kx - 2y}{(K^2 + 2)(m_1 - m_3)} + \frac{\alpha}{2\sqrt{2}} \frac{K(Ky + x)}{(K^2 + 2)(m_2 - m_3)}. \end{aligned} \quad (31)$$

In this case, the lepton mixing matrix U' can still be parameterized in terms of three new Euler's angles θ'_{ij} , which are also a perturbation from the θ_{ij} in the case 1, defined by

$$s'_{13} = -U'_{13} = \Delta U_{13} = -\frac{\alpha y}{2\sqrt{2}B},$$

$$\begin{aligned}
t'_{12} &= -\frac{U'_{12}}{U'_{11}} = -\left[4\alpha B^2 Cx + \alpha C^2(C + \sqrt{C^2 + 8B^2})x + 2BC(C + \sqrt{C^2 + 8B^2})(2C - \alpha y) \right. \\
&\quad \left. + 8B^3(4C + 4\sqrt{C^2 + 8B^2} - \alpha y)\right] / \left\{\sqrt{2}\left[64B^4 + 2C^3(C + \sqrt{C^2 + 8B^2}) \right. \right. \\
&\quad \left. \left. - \alpha BC(C + \sqrt{C^2 + 8B^2})x + 2B^2(12C^2 + 8C\sqrt{C^2 + 8B^2}) + \alpha Cy + \alpha y\sqrt{C^2 + 8B^2}\right]\right\}, \\
t'_{23} &= -\frac{U'_{23}}{U'_{33}} = \frac{4B^2 + \alpha(Bx - Cy)}{4B^2 - \alpha(Bx - Cy)}.
\end{aligned}$$

It is easily shown that our model is consistent because the five experimental constraints on the mixing angles and squared mass differences of neutrinos can be respectively fitted with two Yukawa coupling parameters x , y of the antisextet scalar s , if the VEVs are previously given. Indeed, taking the data in (1) we obtain $\alpha \simeq 0.0692$, $x \simeq 0.0728$, $y \simeq -0.1562$, and $B \simeq -0.0241$, $C = 0.022$, $K = 1.943$, and $t'_{23} = 0.9045$ ($\theta'_{23} \simeq 42.13^\circ$, $\sin^2(2\theta'_{23}) = 0.98999$ satisfying the condition $\sin^2(2\theta'_{23}) > 0.95$ as in (1)). The neutrino masses are explicitly given as $m'_1 \simeq -0.02737\text{eV}$, $m'_2 \simeq -0.02870\text{ eV}$ and $m'_3 \simeq -0.05607\text{ eV}$, which are in a normal ordering. The neutrino mixing matrix then takes the form:

$$U = \begin{pmatrix} 0.8251 & -0.5657 & -0.1585 \\ 0.3302 & 0.6781 & -0.6716 \\ 0.4697 & 0.4888 & 0.7426 \end{pmatrix}. \quad (32)$$

IV. SCALAR POTENTIAL

To be complete, we write the scalar potentials of both the models mentioned. It is also noted that $(\text{Tr}A)(\text{Tr}B) = \text{Tr}(A\text{Tr}B)$ and $V(X \rightarrow X', Y \rightarrow Y', \dots) \equiv V(X, Y, \dots)|_{X=X', Y=Y', \dots}$. The general potential invariant under any subgroup takes the form

$$V_{\text{total}} = V_{\text{tri}} + V_{\text{sext}} + V_{\text{tri-sext}}, \quad (33)$$

where V_{tri} comes from only contributions of $\text{SU}(3)_L$ triplets given as a sum of the following terms:

$$\begin{aligned}
V(\chi) &= \mu_\chi^2 \chi^\dagger \chi + \lambda^\chi (\chi^\dagger \chi)^2, \\
V(\phi) &= V(\chi \rightarrow \phi), \quad V(\phi') = V(\chi \rightarrow \phi'), \quad V(\eta) = V(\chi \rightarrow \eta), \quad V(\eta') = V(\chi \rightarrow \eta'), \\
V(\phi, \chi) &= \lambda_1^{\phi\chi} (\phi^\dagger \phi)(\chi^\dagger \chi) + \lambda_2^{\phi\chi} (\phi^\dagger \chi)(\chi^\dagger \phi), \\
V(\phi', \chi) &= V(\phi \rightarrow \phi', \chi), \quad V(\chi, \eta) = V(\chi, \phi \rightarrow \eta), \quad V(\chi, \eta') = V(\chi, \phi \rightarrow \eta'), \\
V(\phi, \phi') &= V(\phi, \chi \rightarrow \phi') + \lambda_3^{\phi\phi'} (\phi^\dagger \phi')(\phi^\dagger \phi') + \lambda_4^{\phi\phi'} (\phi^\dagger \phi)(\phi'^\dagger \phi), \\
V(\phi, \eta) &= V(\phi, \chi \rightarrow \eta), \quad V(\phi, \eta') = V(\phi, \chi \rightarrow \eta'),
\end{aligned} \quad (34)$$

$$\begin{aligned}
V(\phi', \eta) &= V(\phi \rightarrow \phi', \chi \rightarrow \eta), \quad V(\phi', \eta') = V(\phi \rightarrow \phi', \chi \rightarrow \eta'), \\
V(\eta, \eta') &= V(\phi \rightarrow \eta, \chi \rightarrow \eta') + \lambda_3^{\eta\eta'} (\eta^\dagger \eta') (\eta^\dagger \eta') + \lambda_4^{\eta\eta'} (\eta'^\dagger \eta) (\eta'^\dagger \eta), \\
V_{\chi\phi\phi'\eta\eta'} &= \mu_1 \chi \phi \eta + \mu'_1 \chi \phi' \eta' \\
&\quad + \lambda_1^1 (\phi^\dagger \phi') (\eta^\dagger \eta') + \lambda_1^2 (\phi^\dagger \phi') (\eta'^\dagger \eta) + \lambda_1^3 (\phi^\dagger \eta) (\eta'^\dagger \phi') + \lambda_1^4 (\phi^\dagger \eta') (\eta'^\dagger \phi') + h.c.
\end{aligned} \tag{36}$$

The V_{sext} is given by only $V(s)$:

$$\begin{aligned}
V(s) &= \mu_s^2 \text{Tr}(s^\dagger s) + \lambda_1^s \text{Tr}[(s^\dagger s)_\underline{1} (s^\dagger s)_\underline{1}] + \lambda_2^s \text{Tr}[(s^\dagger s)_{\underline{1}'} (s^\dagger s)_{\underline{1}'}] + \lambda_3^s \text{Tr}[(s^\dagger s)_\underline{2} (s^\dagger s)_\underline{2}] \\
&\quad + \lambda_4^s \text{Tr}(s^\dagger s)_\underline{1} \text{Tr}(s^\dagger s)_\underline{1} + \lambda_5^s \text{Tr}(s^\dagger s)_{\underline{1}'} \text{Tr}(s^\dagger s)_{\underline{1}'} + \lambda_6^s \text{Tr}(s^\dagger s)_\underline{2} \text{Tr}(s^\dagger s)_\underline{2}.
\end{aligned} \tag{37}$$

Next, $V_{\text{tri-sext}}$ is a sum of all the terms connecting both the sectors:

$$\begin{aligned}
V(\phi, s) &= \lambda_1^{\phi s} (\phi^\dagger \phi) \text{Tr}(s^\dagger s)_\underline{1} + \lambda_2^{\phi s} [(\phi^\dagger s^\dagger)(s\phi)]_\underline{1}, \\
V(\phi', s) &= V(\phi \rightarrow \phi', s), \quad V(\chi, s) = V(\phi \rightarrow \chi, s), \\
V(\eta, s) &= V(\phi \rightarrow \eta, s), \quad V(\eta', s) = V(\phi \rightarrow \eta', s), \\
V_{s\chi\phi\phi'\eta\eta'} &= (\lambda'_1 \phi^\dagger \phi' + \lambda'_2 \eta^\dagger \eta') \text{Tr}(s^\dagger s)_{\underline{1}'} + \lambda'_3 [(\phi^\dagger s^\dagger)(s\phi')]_\underline{1} + \lambda'_4 [(\eta^\dagger s^\dagger)(s\eta')]_\underline{1} + h.c.
\end{aligned}$$

To provide the Majorana masses for the neutrinos, the lepton number must be broken. This can be achieved via the scalar potential violating $U(1)_{\mathcal{L}}$, but the other symmetries should be conserved. The \mathcal{L} violating potential is given by

$$\begin{aligned}
\bar{V} &= [\bar{\lambda}_1 \text{Tr}(s^\dagger s)_\underline{1} + \bar{\lambda}_2 \eta^\dagger \chi + \bar{\lambda}_3 \eta^\dagger \eta + \bar{\lambda}_4 \eta'^\dagger \eta' + \bar{\lambda}_5 \eta^\dagger \eta' + \bar{\lambda}_6 \eta'^\dagger \eta + \bar{\lambda}_7 \phi^\dagger \phi \\
&\quad + \bar{\lambda}_8 \phi'^\dagger \phi' + \bar{\lambda}_9 \phi^\dagger \phi' + \bar{\lambda}_{10} \phi'^\dagger \phi] \eta^\dagger \chi + [\bar{\lambda}_{11} \text{Tr}(s^\dagger s)_{\underline{1}'} + \bar{\lambda}_{12} \eta'^\dagger \chi + \bar{\lambda}_{13} \eta^\dagger \eta \\
&\quad + \bar{\lambda}_{14} \eta'^\dagger \eta' + \bar{\lambda}_{15} \eta^\dagger \eta' + \bar{\lambda}_{16} \eta'^\dagger \eta + \bar{\lambda}_{17} \phi^\dagger \phi + \bar{\lambda}_{18} \phi'^\dagger \phi' + \bar{\lambda}_{19} \phi^\dagger \phi' + \bar{\lambda}_{20} \phi'^\dagger \phi]_{\underline{1}'} \eta'^\dagger \chi \\
&\quad + \bar{\lambda}_{21} (\eta^\dagger \phi) (\phi^\dagger \chi) + \bar{\lambda}_{22} (\eta^\dagger \phi')_{\underline{1}'} (\phi'^\dagger \chi)_{\underline{1}'} + \bar{\lambda}_{23} (\eta'^\dagger \phi)_{\underline{1}'} (\phi'^\dagger \chi)_{\underline{1}'} \\
&\quad + \bar{\lambda}_{24} (\eta'^\dagger \phi')_{\underline{1}'} (\phi'^\dagger \chi)_{\underline{1}'} + \bar{\lambda}_{25} (\eta^\dagger s^\dagger)_\underline{2} (s\chi)_\underline{2} + \bar{\lambda}_{26} (\eta'^\dagger s^\dagger)_\underline{2} (s\chi)_\underline{2} + h.c.
\end{aligned} \tag{38}$$

We have not pointed it out, but there must additionally exist the terms in \bar{V} explicitly violating the only S_3 symmetry or both the S_3 and \mathcal{L} -charge. In what follows, most of them will be omitted, only the terms of interest to us are provided.

We now consider the potential V_{tri} . The flavons $\chi, \phi, \phi', \eta, \eta'$ with their VEVs aligned in the same direction (all of them are singlets) are a automatical solution of the minimization conditions for V_{tri} . To explicitly see this, in the system of equations for minimization, we set $v^* = v, v'^* = v', u^* = u, u'^* = u', \text{and } v_\chi^* = v_\chi$. Then the potential minimization conditions for triplets reduces to

$$\frac{\partial V_{\text{tri}}}{\partial \omega} = 4\lambda\chi\omega^3 + 2\left(\mu_\chi^2 + \lambda_1^{\chi\eta}u^2 + \lambda_1^{\chi\eta'}u'^2 + \lambda_1^{\chi\phi}v^2 + \lambda_1^{\chi\phi'}v'^2\right)\omega - \mu_1 uv - \mu'_1 u'v' = 0, \tag{39}$$

$$\begin{aligned} \frac{\partial V_{tri}}{\partial v} &= 4\lambda^\phi v^3 + 2 \left[\mu_\phi^2 + \lambda_1^{\phi\eta} u^2 + \lambda_1^{\phi\eta'} u'^2 + (\lambda_1^{\phi\phi'} + \lambda_2^{\phi\phi'} + \lambda_3^{\phi\phi'} + \lambda_4^{\phi\phi'}) v'^2 + \omega^2 \lambda_1^{\phi\chi} \right] v \\ &+ (\lambda_1^1 + \lambda_1^2) u u' v' - \mu_1 \omega u = 0, \end{aligned} \quad (40)$$

$$\begin{aligned} \frac{\partial V_{tri}}{\partial v'} &= 4\lambda^{\phi'} v'^3 + 2 \left[\mu_{\phi'}^2 + \lambda_1^{\phi'\eta} u^2 + \lambda_1^{\phi'\eta'} u'^2 + (\lambda_1^{\phi\phi'} + \lambda_2^{\phi\phi'} + \lambda_3^{\phi\phi'} + \lambda_4^{\phi\phi'}) v^2 + \omega^2 \lambda_1^{\phi'\chi} \right] v' \\ &+ (\lambda_1^1 + \lambda_1^2) u u' v - \mu_1' \omega u' = 0, \end{aligned} \quad (41)$$

$$\begin{aligned} \frac{\partial V_{tri}}{\partial u} &= 4\lambda^\eta u^3 + 2 \left[\mu_\eta^2 + (\lambda_1^{\eta\eta'} + \lambda_2^{\eta\eta'} + \lambda_3^{\eta\eta'} + \lambda_4^{\eta\eta'}) u'^2 + \lambda_1^{\phi\eta} v^2 + \lambda_1^{\phi'\eta} v'^2 + \omega^2 \lambda_1^{\eta\chi} \right] u \\ &+ (\lambda_1^1 + \lambda_1^2) u' v v' - \mu_1 \omega v = 0, \end{aligned} \quad (42)$$

$$\begin{aligned} \frac{\partial V_{tri}}{\partial u'} &= 4\lambda^{\eta'} u'^3 + 2 \left[\mu_{\eta'}^2 + (\lambda_1^{\eta\eta'} + \lambda_2^{\eta\eta'} + \lambda_3^{\eta\eta'} + \lambda_4^{\eta\eta'}) u^2 + \lambda_1^{\phi\eta'} v^2 + \lambda_1^{\phi'\eta'} v'^2 + \omega^2 \lambda_1^{\eta'\chi} \right] u' \\ &+ (\lambda_1^1 + \lambda_1^2) u v v' - \mu_1' \omega v' = 0. \end{aligned} \quad (43)$$

It is easily to see that the derivatives of V_{tri} with respect to the variables u, u', v, v' shown in (40), (41), (42) and (43) are symmetric with respect to one another. System of equations (39) - (43) always has the solution (u, v, u', v') as expected, even though it is complicated. We also note that the above alignment is only one of the conditions to be imposed to have the desirable results. We have evaluated that Eqs. (40) - (43) have the same structure of solutions. Consequently, to have a simple solution, we can assume that $u = u' = v = v'$. In this case, Eqs. (40) - (43) reduce to a single equation, and system of equations (39) - (43) becomes

$$\frac{\partial V_{tri}}{\partial \omega} = 4\lambda^\chi \omega^3 + 2\omega[\mu_\chi^2 + (2\lambda_1^{\chi\eta} + 2\lambda_1^{\chi\phi})v^2] - 2\mu_1 v^2 = 0, \quad (44)$$

$$\begin{aligned} \frac{\partial V_{tri}}{\partial v} &= 2v \left[2\omega^2(\lambda_1^{\chi\eta} + \lambda_1^{\chi\phi}) + 2(\mu_\eta^2 + \mu_\phi^2) + 2 \left(\lambda_1^1 + \lambda_1^2 + 4\lambda_1^{\phi\eta} + \lambda_1^{\eta\eta'} + \lambda_2^{\eta\eta'} \right. \right. \\ &\left. \left. + \lambda_3^{\eta\eta'} + \lambda_4^{\eta\eta'} + \lambda_1^{\phi\phi'} + \lambda_2^{\phi\phi'} + \lambda_3^{\phi\phi'} + \lambda_4^{\phi\phi'} + 2\lambda^\phi + 2\lambda^\eta \right) v^2 - 2\mu_1 \omega \right] = 0. \end{aligned} \quad (45)$$

This system has the solution

$$\begin{aligned} u &= u' = v' = v = \pm \sqrt{\omega(\mu_\chi^2 + \lambda^\chi \omega^2)} / \sqrt{\mu_1 - 2\omega(\lambda_1^{\chi\eta} + \lambda_1^{\chi\phi})}, \\ \omega &= \frac{\alpha\mu_1}{2(\alpha^2 - \beta\lambda^\chi)} - \frac{\Omega}{3 \times 2^{2/3}(\alpha^2 - \beta\lambda^\chi) \left(\Gamma + \sqrt{\Gamma^2 + 4\Omega^3} \right)^{1/3}} + \frac{\left(\Gamma + \sqrt{\Gamma^2 + 4\Omega^3} \right)^{1/3}}{6 \times 2^{1/3}(\alpha^2 - \beta\lambda^\chi)} \end{aligned}$$

where

$$\Gamma = 54\alpha\beta\mu_1(\lambda^\chi \mu_1^2 + \alpha^2 \mu_\chi^2 - \beta\lambda^\chi \mu_\chi^2) - 108\lambda^\chi \mu_1 \beta \gamma (\alpha^2 - \lambda^\chi \beta), \quad (46)$$

$$\Omega = 6(\alpha^2 - \beta\lambda^\chi)(2\alpha\gamma + \mu_1^2 - \beta\mu_\chi^2) - 9\alpha^2 \mu_1^2, \quad (47)$$

$$\alpha = \lambda_1^{\chi\eta} + \lambda_1^{\chi\phi}, \quad (48)$$

$$\beta = \lambda_1^1 + \lambda_1^2 + 4\lambda_1^{\phi\eta} + \lambda^{\phi\phi'} + \lambda^{\eta\eta'} + 2(\lambda^\eta + \lambda^\phi). \quad (49)$$

$$\lambda^{\phi\phi'} = \lambda_1^{\phi\phi'} + \lambda_2^{\phi\phi'} + \lambda_3^{\phi\phi'} + \lambda_4^{\phi\phi'}, \quad \lambda^{\eta\eta'} = \lambda_1^{\eta\eta'} + \lambda_2^{\eta\eta'} + \lambda_3^{\eta\eta'} + \lambda_4^{\eta\eta'}.$$

We next consider the potential V_{sext} and $V_{tri-sext}$. By imposing the conditions

$$\begin{aligned}\lambda_1^* &= \lambda_1, \lambda_2^* = \lambda_2, v_1^* = v_1, v_2^* = v_2, \Lambda_1^* = \Lambda_1, \Lambda_2^* = \Lambda_2, \\ v^* &= v, v'^* = v', u^* = u, u'^* = u', v_\chi^* = v_\chi, v_\rho^* = v_\rho,\end{aligned}\tag{50}$$

we obtain a system of equations of the potential minimization for anti-sextets:

$$\begin{aligned}\frac{\partial V_1}{\partial \lambda_1} &= 2 \left\{ \lambda_2 \left[\lambda_1^{\chi s} \omega^2 + \mu_s^2 + (\lambda_1^{\eta s} + \lambda_2^{\eta s} + \lambda_3^{\eta s}) u^2 + (\lambda_2' + \lambda_4') u u' + (\lambda_1^{\eta' s} + \lambda_2^{\eta' s} + \lambda_3^{\eta' s}) u'^2 + \lambda_1' v v' \right. \right. \\ &\quad + \lambda_1^{\phi s} v^2 + \lambda_1^{\phi' s} v'^2 + 4 \lambda_4^s \Lambda_1 \Lambda_2 + 2(3 \lambda_1^s + \lambda_2^s + \lambda_3^s + 4 \lambda_4^s) v_1 v_2 \left. \right] + 2 \Lambda_2 (\lambda_1^s - \lambda_2^s + \lambda_3^s) v_1 v_2 \\ &\quad + 2 \Lambda_1 (\lambda_1^s + \lambda_2^s) v_2^2 + 2 \lambda_1 \left[\lambda_6^s \Lambda_2^2 + \lambda_2^2 (2 \lambda_1^s + \lambda_3^s + 2 \lambda_4^s + \lambda_6^s) + (\lambda_1^s - \lambda_2^s + \lambda_3^s + 2 \lambda_6^s) v_2^2 \right] \left. \right\},\end{aligned}\tag{51}$$

$$\begin{aligned}\frac{\partial V_1}{\partial \lambda_2} &= 2 \left\{ \lambda_1 \left[\lambda_1^{\chi s} \omega^2 + \mu_s^2 + (\lambda_1^{\eta s} + \lambda_2^{\eta s} + \lambda_3^{\eta s}) u^2 + (\lambda_2' + \lambda_4') u u' + (\lambda_1^{\eta' s} + \lambda_2^{\eta' s} + \lambda_3^{\eta' s}) u'^2 + \lambda_1' v v' \right. \right. \\ &\quad + \lambda_1^{\phi s} v^2 + \lambda_1^{\phi' s} v'^2 + 4 \lambda_4^s \Lambda_1 \Lambda_2 + 2(3 \lambda_1^s + \lambda_2^s + \lambda_3^s + 4 \lambda_4^s) v_1 v_2 \left. \right] + 2 \Lambda_1 (\lambda_1^s - \lambda_2^s + \lambda_3^s) v_1 v_2 \\ &\quad + 2 \Lambda_2 (\lambda_1^s + \lambda_2^s) v_1^2 + 2 \lambda_2 \left[\lambda_6^s \Lambda_1^2 + \lambda_1^2 (2 \lambda_1^s + \lambda_3^s + 2 \lambda_4^s + \lambda_6^s) + (\lambda_1^s - \lambda_2^s + \lambda_3^s + 2 \lambda_6^s) v_1^2 \right] \left. \right\},\end{aligned}\tag{52}$$

$$\begin{aligned}\frac{\partial V_1}{\partial v_1} &= 2 \left\{ v_2 \left[(2 \lambda_1^{\chi s} + \lambda_2^{\chi s} + \lambda_3^{\chi s}) \omega^2 + 2 \mu_s^2 + (2 \lambda_1^{\eta s} + \lambda_2^{\eta s} + \lambda_3^{\eta s}) u^2 + (2 \lambda_2' + \lambda_4') u u' \right. \right. \\ &\quad + (2 \lambda_1^{\eta' s} + \lambda_2^{\eta' s} + \lambda_3^{\eta' s}) u'^2 + 2 \lambda_1^{\phi s} v^2 + 2 \lambda_1' v v' + 2 \lambda_1^{\phi' s} v'^2 + 2 (\lambda_1 \Lambda_2 + \lambda_2 \Lambda_1) (\lambda_1^s - \lambda_2^s + \lambda_3^s) \\ &\quad + 2 (\lambda_1 \lambda_2 + \Lambda_1 \Lambda_2) (3 \lambda_1^s + \lambda_2^s + \lambda_3^s + 4 \lambda_4^s) \left. \right] + 2 \left[2 \lambda_2 \Lambda_2 (\lambda_1^s + \lambda_2^s) + (\lambda_2^2 + \Lambda_2^2) (\lambda_1^s - \lambda_2^s \right. \\ &\quad + \lambda_3^s + 2 \lambda_6^s) \left. \right] v_1 + 4 (2 \lambda_1^s + \lambda_3^s + 4 \lambda_4^s + 2 \lambda_6^s) v_1 v_2^2 \left. \right\},\end{aligned}\tag{53}$$

$$\begin{aligned}\frac{\partial V_1}{\partial v_2} &= 2 \left\{ v_1 \left[(2 \lambda_1^{\chi s} + \lambda_2^{\chi s} + \lambda_3^{\chi s}) \omega^2 + 2 \mu_s^2 + (2 \lambda_1^{\eta s} + \lambda_2^{\eta s} + \lambda_3^{\eta s}) u^2 + (2 \lambda_2' + \lambda_4') u u' \right. \right. \\ &\quad + (2 \lambda_1^{\eta' s} + \lambda_2^{\eta' s} + \lambda_3^{\eta' s}) u'^2 + 2 \lambda_1^{\phi s} v^2 + 2 \lambda_1' v v' + 2 \lambda_1^{\phi' s} v'^2 + 2 (\lambda_1 \Lambda_2 + \lambda_2 \Lambda_1) (\lambda_1^s - \lambda_2^s + \lambda_3^s) \\ &\quad + 2 (\lambda_1 \lambda_2 + \Lambda_1 \Lambda_2) (3 \lambda_1^s + \lambda_2^s + \lambda_3^s + 4 \lambda_4^s) \left. \right] + 2 \left[2 \lambda_1 \Lambda_1 (\lambda_1^s + \lambda_2^s) + (\lambda_1^2 + \Lambda_1^2) (\lambda_1^s - \lambda_2^s \right. \\ &\quad + \lambda_3^s + 2 \lambda_6^s) \left. \right] v_2 + 4 (2 \lambda_1^s + \lambda_3^s + 4 \lambda_4^s + 2 \lambda_6^s) v_2 v_1^2 \left. \right\},\end{aligned}\tag{54}$$

$$\begin{aligned}\frac{\partial V_1}{\partial \Lambda_1} &= 2 \left\{ \Lambda_2 \left[(\lambda_1^{\chi s} + \lambda_2^{\chi s} + \lambda_3^{\chi s}) \omega^2 + \mu_s^2 + \lambda_1^{\eta s} u^2 + \lambda_2' u u' + \lambda_1^{\eta' s} u'^2 + \lambda_1' v v' + \lambda_1^{\phi s} v^2 + \lambda_1^{\phi' s} v'^2 \right. \right. \\ &\quad + 4 \lambda_4^s \lambda_1 \lambda_2 + 2(3 \lambda_1^s + \lambda_2^s + \lambda_3^s + 4 \lambda_4^s) v_1 v_2 \left. \right] + 2 \lambda_2 (\lambda_1^s - \lambda_2^s + \lambda_3^s) v_1 v_2 + 2 \lambda_1 (\lambda_1^s + \lambda_2^s) v_2^2 \\ &\quad + 2 \Lambda_1 \left[\lambda_6^s \lambda_2^2 + \Lambda_2^2 (2 \lambda_1^s + \lambda_3^s + 2 \lambda_4^s + \lambda_6^s) + (\lambda_1^s - \lambda_2^s + \lambda_3^s + 2 \lambda_6^s) v_2^2 \right] \left. \right\},\end{aligned}\tag{55}$$

$$\begin{aligned}\frac{\partial V_1}{\partial \Lambda_2} &= 2 \left\{ \Lambda_1 \left[(\lambda_1^{\chi s} + \lambda_2^{\chi s} + \lambda_3^{\chi s}) \omega^2 + \mu_s^2 + \lambda_1^{\eta s} u^2 + \lambda_2' u u' + \lambda_1^{\eta' s} u'^2 + \lambda_1' v v' + \lambda_1^{\phi s} v^2 + \lambda_1^{\phi' s} v'^2 \right. \right. \\ &\quad + 4 \lambda_4^s \lambda_1 \lambda_2 + 2(3 \lambda_1^s + \lambda_2^s + \lambda_3^s + 4 \lambda_4^s) v_1 v_2 \left. \right] + 2 \lambda_1 (\lambda_1^s - \lambda_2^s + \lambda_3^s) v_1 v_2 + 2 \lambda_2 (\lambda_1^s + \lambda_2^s) v_1^2 \\ &\quad + 2 \Lambda_2 \left[\lambda_6^s \lambda_1^2 + \Lambda_1^2 (2 \lambda_1^s + \lambda_3^s + 2 \lambda_4^s + \lambda_6^s) + (\lambda_1^s - \lambda_2^s + \lambda_3^s + 2 \lambda_6^s) v_1^2 \right] \left. \right\},\end{aligned}\tag{56}$$

where

$$V_1 = V_{sext} + V_{tri-sext}$$

It is easily to see that Eqs. (52) – (56) take the same form pairwise. This system of equations yields the relations

$$\lambda_1 = \kappa\lambda_2, \quad v_1 = \kappa v_2, \quad \Lambda_1 = \kappa\Lambda_2, \quad (57)$$

with κ is a constant. It means that there are several alignments for VEVs. In this paper, to obtain the desired results, we impose the two directions for breaking $S_3 \rightarrow Z_2$ and $Z_2 \rightarrow \{\text{Identity}\}$ as mentioned, in which $\kappa = 1$ and $\kappa \neq 1$ but approximates to the unit. In the case where $\kappa = 1$ or $\lambda_1 = \lambda_2 = \lambda_s$, $v_1 = v_2 = v_s$ and $\Lambda_1 = \Lambda_2 = \Lambda_s$, system of equations (51) - (56) reduces to a system for the potential minimal consisting of three equations:

$$\lambda_s \left[A_\omega + \mu_s^2 + 2A_s\Lambda_s^2 + 2(A_s + B_s)\lambda_s^2 + A_v + 4(A_s + B_s)v_s^2 \right] + 2B_s\Lambda_s v_s^2 = 0, \quad (58)$$

$$2(A_\omega + B_\omega) + 2\mu_s^2 + A_v + A'_v + 4B_s\lambda_s\Lambda_s + 4(A_s + B_s)(\lambda_s^2 + v_s^2 + \Lambda_s^2) = 0, \quad (59)$$

$$\Lambda_s \left[A_\omega + B_\omega + \mu_s^2 + 2A_s\lambda_s^2 + 2(A_s + B_s)\Lambda_s^2 + A'_v + 4(A_s + B_s)v_s^2 \right] + 2B_s\lambda_s v_s^2 = 0, \quad (60)$$

where

$$\begin{aligned} A_\omega &= \lambda_1^{\chi s} \omega^2, \quad B_\omega = (\lambda_2^{\chi s} + \lambda_3^{\chi s}) \omega^2, \quad A_s = 2\lambda_4^s + \lambda_6^s, \quad B_s = 2\lambda_1^s + \lambda_3^s, \\ A_v &= (\lambda'_1 + \lambda'_2 + \lambda'_4 + \lambda_1^{\phi s} + \lambda_1^{\phi' s} + \lambda_1^{\eta s} + \lambda_2^{\eta s} + \lambda_3^{\eta s} + \lambda_1^{\eta' s} + \lambda_2^{\eta' s} + \lambda_3^{\eta' s}) v^2, \\ A'_v &= (\lambda'_1 + \lambda'_2 + \lambda_1^{\phi s} + \lambda_1^{\phi' s} + \lambda_1^{\eta s} + \lambda_1^{\eta' s}) v^2. \end{aligned}$$

System of equations (58) - (60) always has the solution $(\lambda_s, v_s, \Lambda_s)$ as expected, even though it is complicated. We also note that the above alignment is only one of the conditions to be imposed to have the desired results.

V. GAUGE BOSONS

The covariant derivative of the triplet is given by

$$D_\mu = \partial_\mu - ig \frac{\lambda_a}{2} W_{\mu a} - ig_X X \frac{\lambda_9}{2} B_\mu = \partial_\mu - iP_\mu, \quad (61)$$

where $\lambda_9 = \sqrt{\frac{2}{3}} \text{diag}(1, 1, 1)$ and $\lambda_a (a = 1, 2, \dots, 8)$ are Gell-Mann matrices that satisfy the relations $\text{Tr} \lambda_a \lambda_b = 2\delta_{ab}$ and $\text{Tr} \lambda_9 \lambda_9 = 2$, and X is $U(1)_X$ -charge of Higgs triplets.

We can rewrite P_μ in a convenient form as follows:

$$\frac{g}{2} \begin{pmatrix} W_{\mu 3} + \frac{W_{\mu 8}}{\sqrt{3}} + t\sqrt{\frac{2}{3}}XB_\mu & \sqrt{2}W_\mu'^+ & \sqrt{2}X_\mu'^0, \\ \sqrt{2}W_\mu'^- & -W_{\mu 3} + \frac{W_{\mu 8}}{\sqrt{3}} + t\sqrt{\frac{2}{3}}XB_\mu & \sqrt{2}Y_\mu'^- \\ \sqrt{2}X_\mu'^{0*} & \sqrt{2}Y_\mu'^+ & -\frac{2}{\sqrt{3}}W_{\mu 8} + t\sqrt{\frac{2}{3}}XB_\mu \end{pmatrix}, \quad (62)$$

where we set

$$\begin{aligned} W_\mu'^+ &= \frac{W_{\mu 1} - iW_{\mu 2}}{\sqrt{2}}, \quad X_\mu'^0 = \frac{W_{\mu 4} - iW_{\mu 5}}{\sqrt{2}}, \\ Y_\mu'^- &= \frac{W_{\mu 6} - iW_{\mu 7}}{\sqrt{2}}, \quad W_\mu'^- = (W_\mu'^+)^*, \quad Y_\mu'^+ = (Y_\mu'^-)^*, \end{aligned} \quad (63)$$

and $t = g_X/g$. We note that W_4 and W_5 are respectively purely real and imaginary parts of X^0 and X^{0*} . The covariant derivative for the antisextet with a VEV part is [22, 23]

$$D_\mu \langle s_i \rangle = \frac{ig}{2} \{ W_\mu^a \lambda_a^* \langle s_i \rangle + \langle s_i \rangle W_\mu^a \lambda_a^{*T} \} - ig_X T_9 X B_\mu \langle s_i \rangle. \quad (64)$$

The covariant derivative (64) acting on the antisextet VEVs is given by

$$\begin{aligned} [D_\mu \langle s_i \rangle]_{11} &= ig \left(\lambda_i W_{\mu 3} + \frac{\lambda_i}{\sqrt{3}} W_{\mu 8} + \sqrt{\frac{2}{3}} \frac{1}{3} t \lambda_i B_\mu + \sqrt{2} v_i X_\mu'^{0*} \right), \\ [D_\mu \langle s_i \rangle]_{12} &= \frac{ig}{\sqrt{2}} \left(\lambda_i W_\mu'^+ + v_i Y_\mu'^+ \right), \\ [D_\mu \langle s_i \rangle]_{13} &= \frac{ig}{2} \left(v_i W_{\mu 3} - \frac{v_i}{\sqrt{3}} W_{\mu 8} + \frac{2}{3} \sqrt{\frac{2}{3}} t v_i B_\mu + \sqrt{2} \lambda_i X_\mu'^0 + \sqrt{2} \Lambda_i X_\mu'^{0*} \right), \\ [D_\mu \langle s_i \rangle]_{21} &= [D_\mu \langle s_i \rangle]_{12}, \\ [D_\mu \langle s_i \rangle]_{22} &= 0, \\ [D_\mu \langle s_i \rangle]_{23} &= \frac{ig}{\sqrt{2}} \left(v_i W_\mu'^+ + \Lambda_i Y_\mu'^+ \right), \\ [D_\mu \langle s_i \rangle]_{31} &= [D_\mu \langle s_i \rangle]_{13}, \\ [D_\mu \langle s_i \rangle]_{32} &= [D_\mu \langle s_i \rangle]_{23}, \\ [D_\mu \langle s_i \rangle]_{33} &= ig \left(-\frac{2}{\sqrt{3}} \Lambda_i W_{\mu 8} + \sqrt{\frac{2}{3}} \frac{1}{3} t \Lambda_i B_\mu + \sqrt{2} v_i X_\mu'^0 \right). \end{aligned}$$

The masses of gauge bosons in this model are defined as

$$\begin{aligned} \mathcal{L}_{mass}^{GB} &= (D_\mu \langle \phi \rangle)^+ (D^\mu \langle \phi \rangle) + (D_\mu \langle \phi' \rangle)^+ (D^\mu \langle \phi' \rangle) + (D_\mu \langle \chi \rangle)^+ (D^\mu \langle \chi \rangle) \\ &+ (D_\mu \langle \eta \rangle)^+ (D^\mu \langle \eta \rangle) + (D_\mu \langle \eta' \rangle)^+ (D^\mu \langle \eta' \rangle) \\ &+ \text{Tr}[(D_\mu \langle s_1 \rangle)^+ (D^\mu \langle s_1 \rangle)] + \text{Tr}[(D_\mu \langle s_2 \rangle)^+ (D^\mu \langle s_2 \rangle)]. \end{aligned} \quad (65)$$

Substituting the Higgs VEVs of the model in (65) yields

$$\mathcal{L}_{mass}^{GB} = \frac{v^2}{324} \left[81g^2(W_{\mu 1}^2 + W_{\mu 2}^2) + 81g^2(W_{\mu 6}^2 + W_{\mu 7}^2) + (-9gW_{\mu 3} + 3\sqrt{3}gW_{\mu 8} + 2\sqrt{6}g_X B_\mu)^2 \right]$$

$$\begin{aligned}
& + \frac{v'^2}{324} \left[81g^2(W_{\mu 1}^2 + W_{\mu 2}^2) + 81g^2(W_{\mu 6}^2 + W_{\mu 7}^2) + (-9gW_{\mu 3} + 3\sqrt{3}gW_{\mu 8} + 2\sqrt{6}g_X B_\mu)^2 \right] \\
& + \frac{\omega^2}{108} \left[27g^2(W_{\mu 4}^2 + W_{\mu 5}^2) + 27g^2(W_{\mu 6}^2 + W_{\mu 7}^2) + 36g^2W_{\mu 8}^2 + 12\sqrt{2}gg_X W_{\mu 8}B_\mu + 2g_X^2 B_\mu^2 \right] \\
& + \frac{u^2}{324} \left[81g^2(W_{\mu 1}^2 + W_{\mu 2}^2) + 81g^2(W_{\mu 4}^2 + W_{\mu 5}^2) + (-9gW_{\mu 3} - 3\sqrt{3}gW_{\mu 8} + \sqrt{6}g_X B_\mu)^2 \right] \\
& + \frac{u'^2}{324} \left[81g^2(W_{\mu 1}^2 + W_{\mu 2}^2) + 81g^2(W_{\mu 4}^2 + W_{\mu 5}^2) + (-9gW_{\mu 3} - 3\sqrt{3}gW_{\mu 8} + \sqrt{6}g_X B_\mu)^2 \right] \\
& + \frac{g^2}{6} \left[2(\Lambda_1 v_1 + \Lambda_2 v_2) \left(3W_{\mu 3}W_{\mu 4} + 3W_{\mu 1}W_{\mu 6} - 3W_{\mu 2}W_{\mu 7} - 5\sqrt{3}W_{\mu 4}W_{\mu 8} \right) \right. \\
& + 3(v_1^2 + v_2^2 + \lambda_1^2 + \lambda_2^2)W_{\mu 1}^2 + 3(v_1^2 + v_2^2 + \lambda_1^2 + \lambda_2^2)W_{\mu 2}^2 + 3(v_1^2 + v_2^2 + 2\lambda_1^2 + 2\lambda_2^2)W_{\mu 3}^2 \\
& + 3(4v_1^2 + 4v_2^2 + \lambda_1^2 + \lambda_2^2 + \Lambda_1^2 + \Lambda_2^2 + 2\Lambda_1\lambda_1 + 2\Lambda_2\lambda_2)W_{\mu 4}^2 \\
& + 3(4v_1^2 + 4v_2^2 + \lambda_1^2 + \lambda_2^2 + \Lambda_1^2 + \Lambda_2^2 - 2\Lambda_1\lambda_1 - 2\Lambda_2\lambda_2)W_{\mu 5}^2 \\
& + 3(v_1^2 + v_2^2 + \Lambda_1^2 + \Lambda_2^2)W_{\mu 6}^2 + 3(v_1^2 + v_2^2 + \Lambda_1^2 + \Lambda_2^2)W_{\mu 7}^2 \\
& + 2\sqrt{3}(-v_1^2 - v_2^2 + 2\lambda_1^2 + 2\lambda_2^2)W_{\mu 3}W_{\mu 8} + (v_1^2 + v_2^2 + 2\lambda_1^2 + 2\lambda_2^2 + 8\Lambda_1^2 + 8\Lambda_2^2)W_{\mu 8}^2 \\
& + 18(\lambda_1 v_1 + \lambda_2 v_2)W_{\mu 3}W_{\mu 4} + 6(\lambda_1 v_1 + \lambda_2 v_2)W_{\mu 1}W_{\mu 6} - 6(\lambda_1 v_1 + \lambda_2 v_2)W_{\mu 2}W_{\mu 7} \\
& \left. + 2\sqrt{3}(\lambda_1 v_1 + \lambda_2 v_2)W_{\mu 4}W_{\mu 8} \right] \\
& + \frac{2}{27}t^2g^2(\lambda_1^2 + \lambda_2^2 + \Lambda_1^2 + \Lambda_2^2 + 2v_1^2 + 2v_2^2)B_\mu^2 - \frac{2}{3}\sqrt{\frac{2}{3}}tg^2(\lambda_1^2 + \lambda_2^2 + v_1^2 + v_2^2)W_{\mu 3}B_\mu \\
& - \frac{4}{3}\sqrt{\frac{2}{3}}tg^2[(\lambda_1 + \Lambda_1)v_1 + (\lambda_2 + \Lambda_2)v_2]W_{\mu 4}B_\mu \\
& - \frac{2\sqrt{2}}{9}tg^2(\lambda_1^2 + \lambda_2^2 - v_1^2 - v_2^2 - 2\Lambda_1^2 - 2\Lambda_2^2)W_{\mu 8}B_\mu.
\end{aligned} \tag{66}$$

We can split \mathcal{L}_{mass}^{GB} in (66) as

$$\mathcal{L}_{mass}^{GB} = \mathcal{L}_{mass}^{W_5} + \mathcal{L}_{mix}^{CGB} + \mathcal{L}_{mix}^{NGB}, \tag{67}$$

where $\mathcal{L}_{mass}^{W_5}$ is the Lagrangian part of the imaginary part W_5 . This boson is decoupled with its mass given by

$$\mathcal{L}^{W_5} = \frac{g^2}{4} \left(\omega^2 + u^2 + u'^2 + 8v_1^2 + 8v_2^2 + 2\lambda_1^2 + 2\lambda_2^2 + 2\Lambda_1^2 + 2\Lambda_2^2 - 4\Lambda_1\lambda_1 - 4\Lambda_2\lambda_2 \right) W_{\mu 5}^2.$$

Hence,

$$M_{W_5}^2 = \frac{g^2}{2} \left(\omega^2 + u^2 + u'^2 + 8v_1^2 + 8v_2^2 + 2\lambda_1^2 + 2\lambda_2^2 + 2\Lambda_1^2 + 2\Lambda_2^2 - 4\Lambda_1\lambda_1 - 4\Lambda_2\lambda_2 \right). \tag{68}$$

In the limit $\lambda_1, \lambda_2, v_1, v_2 \rightarrow 0$, we have

$$M_{W_5}^2 = \frac{g^2}{2} \left(\omega^2 + u^2 + u'^2 + 2\Lambda_1^2 + 2\Lambda_2^2 \right). \tag{69}$$

Next,

$$\begin{aligned}
\mathcal{L}_{mix}^{CGB} = & \frac{g^2}{4} \left[v^2 + v'^2 + u^2 + u'^2 + 2(v_1^2 + v_2^2 + \lambda_1^2 + \lambda_2^2) \right] (W_{\mu 1}^2 + W_{\mu 2}^2) \\
& + \frac{g^2}{4} \left[v^2 + v'^2 + \omega^2 + 2(v_1^2 + v_2^2 + \Lambda_1^2 + \Lambda_2^2) \right] (W_{\mu 6}^2 + W_{\mu 7}^2) \\
& + g^2 (\Lambda_1 v_1 + \lambda_1 v_1 + \Lambda_2 v_2 + \lambda_2 v_2) (W_{\mu 1} W_{\mu 6} - W_{\mu 2} W_{\mu 7})
\end{aligned} \tag{70}$$

is the Lagrangian part of the charged gauge bosons W and Y , which can be rewritten in matrix form as

$$\mathcal{L}_{mix}^{CGB} = \frac{g^2}{4} (W_{\mu}^{'-} \ Y_{\mu}^{'-}) M_{WY}^2 (W_{\mu}^{'+} \ Y_{\mu}^{'+})^T,$$

where

$$M_{WY}^2 = 2 \begin{pmatrix} v^2 + v'^2 + u^2 + u'^2 + 2(v_1^2 + v_2^2 + \lambda_1^2 + \lambda_2^2) & 2(\Lambda_1 v_1 + \lambda_1 v_1 + \Lambda_2 v_2 + \lambda_2 v_2) \\ 2(\Lambda_1 v_1 + \lambda_1 v_1 + \Lambda_2 v_2 + \lambda_2 v_2) & v^2 + v'^2 + \omega^2 + 2(v_1^2 + v_2^2 + \Lambda_1^2 + \Lambda_2^2) \end{pmatrix}. \tag{71}$$

The matrix M_{WY}^2 in (71) can be diagonalised as

$$U_2^T M_{WY}^2 U_2 = \text{diag}(M_W^2, M_Y^2),$$

where

$$\begin{aligned}
M_W^2 &= \frac{g^2}{4} \left\{ 2(\lambda_1^2 + \lambda_2^2 + 2v_1^2 + 2v_2^2 + \Lambda_1^2 + \Lambda_2^2) + \omega^2 + u^2 + u'^2 + 2(v^2 + v'^2) - \sqrt{\Gamma} \right\}, \\
M_Y^2 &= \frac{g^2}{4} \left\{ 2(\lambda_1^2 + \lambda_2^2 + 2v_1^2 + 2v_2^2 + \Lambda_1^2 + \Lambda_2^2) + \omega^2 + u^2 + u'^2 + 2(v^2 + v'^2) + \sqrt{\Gamma} \right\},
\end{aligned} \tag{72}$$

with

$$\begin{aligned}
\Gamma &= 4\lambda_1^4 + 4\Lambda_1^4 + (2\lambda_2^2 - 2\Lambda_2^2 - \omega^2 + u^2 + u'^2)^2 - 4\lambda_1^2 (2\Lambda_1^2 - 2\lambda_2^2 + 2\Lambda_2^2 + \omega^2 \\
&\quad - u^2 - u'^2 - 4v_1^2) - 4\Lambda_1^2 (2\lambda_2^2 - 2\Lambda_2^2 - \omega^2 + u^2 + u'^2 - 4v_1^2) + 32\Lambda_1 (\lambda_2 + \Lambda_2) v_1 v_2 \\
&\quad + 16(\lambda_2 + \Lambda_2)^2 v_2^2 + 32\lambda_1 v_1 (\Lambda_1 v_1 + \lambda_2 v_2 + \Lambda_2 v_2).
\end{aligned} \tag{73}$$

In our model, the following limits are often used:

$$\lambda_{1,2}^2, v_{1,2}^2 \ll u^2, u'^2, v^2, v'^2, \tag{74}$$

$$u^2, u'^2, v^2, v'^2 \ll \omega^2 \sim \Lambda_{1,2}^2. \tag{75}$$

With the help of (74), Γ in (73) becomes

$$\Gamma \simeq (2\Lambda_1^2 + 2\Lambda_2^2 + \omega^2 - u^2 - u'^2) + \frac{16\Lambda_1 \Lambda_2 v_1 v_2 + 8\Lambda_2^2 v_2^2}{2\Lambda_1^2 + 2\Lambda_2^2 + \omega^2 - u^2 - u'^2}. \tag{76}$$

It then follows that

$$M_W^2 \simeq \frac{g^2}{2} (u^2 + u'^2 + v^2 + v'^2) - \frac{g^2}{2} \Delta_{M_w^2}, \quad (77)$$

with

$$\Delta_{M_w^2} = \frac{4(2\Lambda_1\Lambda_2v_1v_2 + \Lambda_2^2v_2^2)}{2\Lambda_1^2 + 2\Lambda_2^2 + \omega^2 - u^2 - u'^2}. \quad (78)$$

The corresponding eigenstates are arranged into the charged gauge boson mixing matrix

$$U_2 = \begin{pmatrix} \frac{\mathcal{R}}{\sqrt{\mathcal{R}^2+1}} & -\frac{1}{\sqrt{\mathcal{R}^2+1}} \\ \frac{1}{\sqrt{\mathcal{R}^2+1}} & \frac{\mathcal{R}}{\sqrt{\mathcal{R}^2+1}} \end{pmatrix} \equiv \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix},$$

where

$$\mathcal{R} = \frac{2\lambda_1^2 - 2\Lambda_1^2 + 2\lambda_2^2 - 2\Lambda_2^2 - \omega^2 + u^2 + u'^2 - \sqrt{\Gamma}}{4(\lambda_1 + \Lambda_1)v_1 + 4(\lambda_2 + \Lambda_2)v_2}.$$

The physical charged gauge bosons is defined as

$$\begin{aligned} W_\mu^- &= \cos \theta W_\mu'^- + \sin \theta Y_\mu'^-, \\ Y_\mu^- &= -\sin \theta W_\mu'^- + \cos \theta Y_\mu'^-. \end{aligned}$$

The mixing angle θ is given by

$$\begin{aligned} \tan \theta &= \frac{1}{\mathcal{R}} = \frac{4(\lambda_1 + \Lambda_1)v_1 + 4(\lambda_2 + \Lambda_2)v_2}{2\lambda_1^2 - 2\Lambda_1^2 + 2\lambda_2^2 - 2\Lambda_2^2 - \omega^2 + u^2 + u'^2 - \sqrt{\Gamma}} \\ &\simeq \frac{4\Lambda_1v_1 + 4\Lambda_2v_2}{-2\Lambda_1^2 - 2\Lambda_2^2 - \omega^2 - 2(\Lambda_1^2 + \Lambda_2^2)} \sim \frac{v_i}{\Lambda_i}, \quad (i = 1, 2) \end{aligned} \quad (79)$$

We note that in the limit $v_{1,2} \rightarrow 0$ the mixing angle θ tends to zero, $\Gamma = 2\Lambda_1^2 + 2\Lambda_2^2 + \omega^2 - u^2 - u'^2$, and we have

$$\begin{aligned} M_W^2 &= \frac{g^2}{2} (u^2 + u'^2 + v^2 + v'^2), \\ M_Y^2 &= \frac{g^2}{2} (2\Lambda_1^2 + 2\Lambda_2^2 + \omega^2 + v^2 + v'^2). \end{aligned} \quad (80)$$

There is a mixing among the neutral gauge bosons W_3, W_8, B and W_4 . The mass Lagrangian in this case has the form

$$\begin{aligned} \mathcal{L}_{mix}^{NGB} &= \frac{v^2}{324} \left(81g^2W_{\mu 3}^2 + 27g^2W_{\mu 8}^2 + 24g_X^2B_\mu^2 - 54\sqrt{3}g^2W_{\mu 3}W_{\mu 8} - 36\sqrt{6}gg_XW_{\mu 3}B_\mu \right. \\ &\quad \left. + 36\sqrt{2}gg_XW_{\mu 8}B_\mu \right) + \frac{v'^2}{324} \left(81g^2W_{\mu 3}^2 + 27g^2W_{\mu 8}^2 + 24g_X^2B_\mu^2 - 54\sqrt{3}g^2W_{\mu 3}W_{\mu 8} \right. \\ &\quad \left. - 36\sqrt{6}gg_XW_{\mu 3}B_\mu + 36\sqrt{2}gg_XW_{\mu 8}B_\mu \right) + \frac{\omega^2}{108} \left(27g^2W_{\mu 4}^2 + 36g^2W_{\mu 8}^2 \right. \end{aligned}$$

$$\begin{aligned}
& + 12\sqrt{2}gg_x W_{\mu 8} B_\mu + 2g_X^2 B_\mu^2 \Big) + \frac{u^2}{324} \Big(81g^2 W_{\mu 4}^2 + 81g^2 W_{\mu 3}^2 + 27g^2 W_{\mu 8}^2 + 6g_X^2 B_\mu^2 \\
& + 54\sqrt{3}g^2 W_{\mu 3} W_{\mu 8} - 18\sqrt{6}gg_X W_{\mu 3} B_\mu - 18\sqrt{2}W_{\mu 8} B_\mu \Big) + \frac{u'^2}{324} \Big(81g^2 W_{\mu 4}^2 \\
& + 81g^2 W_{\mu 3}^2 + 27g^2 W_{\mu 8}^2 + 6g_X^2 B_\mu^2 + 54\sqrt{3}g^2 W_{\mu 3} W_{\mu 8} - 18\sqrt{6}gg_X W_{\mu 3} B_\mu - 18\sqrt{2}W_{\mu 8} B_\mu \Big) \\
& + \frac{g^2}{6} \Big[2(\Lambda_1 v_1 + \Lambda_2 v_2) \Big(3W_{\mu 3} W_{\mu 4} - 5\sqrt{3}W_{\mu 4} W_{\mu 8} \Big) + 3(v_1^2 + v_2^2 + 2\lambda_1^2 + 2\lambda_2^2) W_{\mu 3}^2 \\
& + 3(4v_1^2 + 4v_2^2 + \lambda_1^2 + \lambda_2^2 + \Lambda_1^2 + \Lambda_2^2 + 2\Lambda_1 \lambda_1 + 2\Lambda_2 \lambda_2) W_{\mu 4}^2 \\
& + 2\sqrt{3}(-v_1^2 - v_2^2 + 2\lambda_1^2 + 2\lambda_2^2) W_{\mu 3} W_{\mu 8} + (v_1^2 + v_2^2 + 2\lambda_1^2 + 2\lambda_2^2 + 8\Lambda_1^2 + 8\Lambda_2^2) W_{\mu 8}^2 \\
& + 18(\lambda_1 v_1 + \lambda_2 v_2) W_{\mu 3} W_{\mu 4} + 2\sqrt{3}(\lambda_1 v_1 + \lambda_2 v_2) W_{\mu 4} W_{\mu 8} \Big] \\
& + \frac{2}{27} t^2 g^2 (\lambda_1^2 + \lambda_2^2 + \Lambda_1^2 + \Lambda_2^2 + 2v_1^2 + 2v_2^2) B_\mu^2 - \frac{2}{3} \sqrt{\frac{2}{3}} t g^2 (\lambda_1^2 + \lambda_2^2 + v_1^2 + v_2^2) W_{\mu 3} B_\mu \\
& - \frac{4}{3} \sqrt{\frac{2}{3}} t g^2 [(\lambda_1 + \Lambda_1) v_1 + (\lambda_2 + \Lambda_2) v_2] W_{\mu 4} B_\mu \\
& - \frac{2\sqrt{2}}{9} t g^2 (\lambda_1^2 + \lambda_2^2 - v_1^2 - v_2^2 - 2\Lambda_1^2 - 2\Lambda_2^2) W_{\mu 8} B_\mu.
\end{aligned} \tag{81}$$

In the basis of $(W_{\mu 3}, W_{\mu 8}, B_\mu, W_{\mu 4})$, \mathcal{L}_{mix}^{NGB} can be rewritten in matrix form:

$$\mathcal{L}_{mix}^{NGB} \equiv \frac{1}{2} V^T M^2 V,$$

where

$$\begin{aligned}
V^T &= (W_{\mu 3}, W_{\mu 8}, B_\mu, W_{\mu 4}), \\
M^2 &= \frac{g^2}{4} \begin{pmatrix} M_{11}^2 & M_{12}^2 & M_{13}^2 & M_{14}^2 \\ M_{12}^2 & M_{22}^2 & M_{23}^2 & M_{24}^2 \\ M_{13}^2 & M_{23}^2 & M_{33}^2 & M_{34}^2 \\ M_{14}^2 & M_{24}^2 & M_{34}^2 & M_{44}^2 \end{pmatrix},
\end{aligned} \tag{82}$$

with

$$\begin{aligned}
M_{11}^2 &= 2(v^2 + v'^2 + u^2 + u'^2 + 2v_1^2 + 2v_2^2 + 4\lambda_1^2 + 4\lambda_2^2), \\
M_{12}^2 &= -\frac{2\sqrt{3}}{3} (v^2 + v'^2 - u^2 - u'^2 + 2v_1^2 + 2v_2^2 - 4\lambda_1^2 - 4\lambda_2^2), \\
M_{13}^2 &= -\frac{2}{3} \sqrt{\frac{2}{3}} t (2v^2 + 2v'^2 + u^2 + u'^2 + 4\lambda_1^2 + 4\lambda_2^2 + 4v_1^2 + 4v_2^2), \\
M_{14}^2 &= 4(\Lambda_1 v_1 + \Lambda_2 v_2) + 12(\lambda_1 v_1 + \lambda_2 v_2), \\
M_{22}^2 &= \frac{2}{3} (v^2 + v'^2 + 4\omega^2 + u^2 + u'^2 + 2v_1^2 + 2v_2^2 + 4\lambda_1^2 + 4\lambda_2^2 + 16\Lambda_1^2 + 16\Lambda_2^2), \\
M_{23}^2 &= \frac{2\sqrt{2}t}{9} (2v^2 + 2v'^2 + 2\omega^2 - u^2 - u'^2 - 4\lambda_1^2 - 4\lambda_2^2 + 4v_1^2 + 4v_2^2 + 8\Lambda_1^2 + 8\Lambda_2^2), \\
M_{24}^2 &= \frac{4}{\sqrt{3}} [\lambda_1 v_1 + \lambda_2 v_2 - 5(\Lambda_1 v_1 + \Lambda_2 v_2)],
\end{aligned}$$

$$\begin{aligned}
M_{33}^2 &= \frac{4t^2}{27} \left(4v^2 + 4v'^2 + \omega^2 + u^2 + u'^2 + 4\lambda_1^2 + 4\lambda_2^2 + 4\Lambda_1^2 + 4\Lambda_2^2 + 8v_1^2 + 8v_2^2 \right), \\
M_{34}^2 &= -\frac{16}{3} \sqrt{\frac{2}{3}} t (\lambda_1 v_1 + \Lambda_1 v_1 + \lambda_2 v_2 + \Lambda_2 v_2), \\
M_{44}^2 &= 2(\omega^2 + u^2 + u'^2 + 8v_1^2 + 8v_2^2 + 2\lambda_1^2 + 2\lambda_2^2 + 2\Lambda_1^2 + 2\Lambda_2^2 + 4\Lambda_1 \lambda_1 + 4\Lambda_2 \lambda_2).
\end{aligned} \tag{83}$$

The matrix M^2 in (82) with the elements in (83) has one exact eigenvalue, which is identified with the photon mass,

$$M_\gamma^2 = 0. \tag{84}$$

The corresponding eigenvector of M_γ^2 is

$$A_\mu = \begin{pmatrix} \frac{\sqrt{3}t}{\sqrt{4t^2+18}} & -\frac{t}{\sqrt{4t^2+18}} & \frac{3\sqrt{2}}{\sqrt{4t^2+18}} & 0 \end{pmatrix}^T. \tag{85}$$

We note that in the limit $\lambda_{1,2}, v_{1,2} \rightarrow 0$, $M_{14}^2 = M_{24}^2 = M_{34}^2 = 0$ and W_4 does not mix with $W_{3\mu}, W_{8\mu}$ and B_μ . In the general case $\lambda_{1,2}, v_{1,2} \neq 0$, the mass matrix in (82) contains one exact eigenvalues as in (84) with the corresponding eigenstate defined in (85).

The diagonalization of the mass matrix M^2 in (82) is done in two steps. In the first step, the basis $(W_{\mu 3}, W_{\mu 8}, B'_\mu, W_{4\mu})$ is transformed into the basis $(A_\mu, Z_\mu, Z'_\mu, W_{4\mu})$ by the matrix

$$U_{NGB} = \begin{pmatrix} s_W & -c_W & 0 & 0 \\ -\frac{c_W t_W}{\sqrt{3}} & -\frac{s_W t_W}{\sqrt{3}} & \sqrt{1 - \frac{t_W^2}{3}} & 0 \\ c_W \sqrt{1 - \frac{t_W^2}{3}} & s_W \sqrt{1 - \frac{t_W^2}{3}} & \frac{t_W}{\sqrt{3}} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \tag{86}$$

The eigenstates are defined as

$$\begin{aligned}
A_\mu &= s_W W_{3\mu} + c_W \left(-\frac{t_W}{\sqrt{3}} W_{8\mu} + \sqrt{1 - \frac{t_W^2}{3}} B_\mu \right), \\
Z_\mu &= -c_W W_{3\mu} + s_W \left(-\frac{t_W}{\sqrt{3}} W_{8\mu} + \sqrt{1 - \frac{t_W^2}{3}} B_\mu \right), \\
Z'_\mu &= \sqrt{1 - \frac{t_W^2}{3}} W_{8\mu} + \frac{t_W}{\sqrt{3}} B_\mu.
\end{aligned} \tag{87}$$

To obtain (86) and (87) we used the continuation of the $SU(3)_L$ gauge coupling constant g to the spontaneous symmetry breaking point, where

$$t = \frac{3\sqrt{2}s_W}{\sqrt{3 - 4s_W^2}}. \tag{88}$$

In this basis, the mass matrix M^2 in (82) becomes

$$M'^2 = U_{NGB}^+ M^2 U_{NGB} = \frac{g^2}{4} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & M'_{22} & M'_{23} & M'_{24} \\ 0 & M'_{23} & M'_{33} & M'_{34} \\ 0 & M'_{24} & M'_{34} & M'_{44} \end{pmatrix}, \quad (89)$$

where

$$\begin{aligned} M'_{22} &= \frac{4(2t^2 + 9)}{t^2 + 18} (u^2 + u'^2 + v^2 + v'^2 + 4\lambda_1^2 + 4\lambda_2^2 + 2v_1^2 + 2v_2^2) \\ &= \frac{2}{c_W^2} (u^2 + u'^2 + v^2 + v'^2 + 4\lambda_1^2 + 4\lambda_2^2 + 2v_1^2 + 2v_2^2), \\ M'_{23} &= \frac{4}{3\sqrt{3}} \frac{\sqrt{2t^2 + 9}}{(t^2 + 18)} \left[(t^2 - 9)(4\lambda_1^2 + 4\lambda_2^2 + u^2 + u'^2) + (2t^2 + 9)(v^2 + v'^2 + 2v_1^2 + 2v_2^2) \right] \\ &= \frac{2[(1 - 2c_W^2)(u^2 + u'^2 + 4\lambda_1^2 + 4\lambda_2^2) + v^2 + v'^2 + v_1^2 + v_2^2]\sqrt{\alpha_0}}{c_W^2}, \\ M'_{24} &= -4\sqrt{2} \sqrt{\frac{2t^2 + 9}{t^2 + 18}} (\Lambda_1 v_1 + 3\lambda_1 v_1 + \Lambda_2 v_2 + 3\lambda_2 v_2) \\ &= -\frac{4}{c_W} (\Lambda_1 v_1 + \Lambda_2 v_2 + 3\lambda_1 v_1 + 3\lambda_2 v_2), \\ M'_{33} &= \frac{4}{27(t^2 + 18)} \left[4\lambda_1^2(t^2 - 9)^2 + 4\lambda_1^2(t^2 + 18)^2 + 81(4\lambda_2^2 + 16\Lambda_2^2 + 4\omega^2 + u^2 + u'^2 \right. \\ &\quad + v^2 + v'^2 + 2v_1^2 + 2v_2^2) + 18t^2(8\Lambda_2^2 + 2\omega^2 - u^2 - u'^2 + 2v^2 + 2v'^2 + 4v_1^2 + 4v_2^2) \\ &\quad + 4\lambda_2^2 t^2(t^2 - 18) + t^4(4\Lambda_2^2 + \omega^2 + u^2 + u'^2 + 4v^2 + 4v'^2 + 8v_1^2 + 8v_2^2) \left. \right] \\ &= 32(\Lambda_1^2 + \Lambda_2^2)c_W^2\alpha_0 + 8\omega^2 c_W^2\alpha_0 + \frac{2}{c_W^2}(v^2 + v'^2 + 2v_1^2 + 2v_2^2)\alpha_0 \\ &\quad + \frac{2}{c_W^2}(2c_W^2 - 1)^2(u^2 + u'^2)\alpha_0 + \frac{8(2c_W^2 - 1)^2}{c_W^2}(\lambda_1^2 + \lambda_2^2)\alpha_0, \\ M'_{34} &= -\frac{4\sqrt{2}}{3\sqrt{3}} \frac{1}{\sqrt{t^2 + 18}} \left[(4t^2 - 9)(\lambda_1 v_1 + \lambda_2 v_2) + (4t^2 + 45)(\Lambda_1 v_1 + \Lambda_2 v_2) \right] \\ &= -\frac{4\sqrt{\alpha_0}}{c_W} \left[x_0(\Lambda_1 v_1 + \Lambda_2 v_2) + \left(2 - \frac{1}{\alpha_0}\right)(\lambda_1 v_1 + \lambda_2 v_2) \right], \\ M'_{44} &= 2 \left[2(\lambda_1 + \Lambda_1)^2 + 2(\lambda_2 + \Lambda_2)^2 + \omega^2 + u^2 + u'^2 + 8v_1^2 + 8v_2^2 \right] \\ &= 2 \left(u^2 + u'^2 + \omega^2 + 2\lambda_1^2 + 2\lambda_2^2 + 2\Lambda_1^2 + 2\Lambda_2^2 + 4\lambda_1\Lambda_1 + 4\lambda_2\Lambda_2 + 8v_1^2 + 8v_2^2 \right). \end{aligned} \quad (90)$$

In the approximation $\lambda_{1,2}^2, v_{1,2}^2 \ll \Lambda_{1,2}^2 \sim \omega^2$, we have

$$\begin{aligned} M'_{22} &= \frac{2}{c_W^2} (u^2 + u'^2 + v^2 + v'^2), \\ M'_{23} &= \frac{2[(1 - 2c_W^2)(u^2 + u'^2) + v^2 + v'^2]\sqrt{\alpha_0}}{c_W^2}, \end{aligned}$$

$$\begin{aligned}
M_{24}'^2 &= -\frac{4}{c_W} (\Lambda_1 v_1 + \Lambda_2 v_2), \\
M_{33}'^2 &= 32(\Lambda_1^2 + \Lambda_2^2) c_W^2 \alpha_0 + 8\omega^2 c_W^2 \alpha_0 + \frac{2}{c_W^2} (v^2 + v'^2) \alpha_0 + \frac{2}{c_W^2} (2c_W^2 - 1)^2 (u^2 + u'^2) \alpha_0, \\
M_{34}'^2 &= -\frac{4x_0 \sqrt{\alpha}}{c_W} (\Lambda_1 v_1 + \Lambda_2 v_2), \\
M_{44}'^2 &= 2(u^2 + u'^2 + \omega^2 + 2\Lambda_1^2 + 2\Lambda_2^2 + 4\lambda_1 \Lambda_1 + 4\lambda_2 \Lambda_2),
\end{aligned} \tag{91}$$

with

$$\begin{aligned}
s_W &= \sin \theta_W, \quad c_W = \cos \theta_W, \quad t_W = \tan \theta_W, \\
x_0 &= 4c_W^2 + 1, \quad \alpha_0 = (4c_W^2 - 1)^{-1}.
\end{aligned} \tag{92}$$

From (89), there exist mixings between Z_μ, Z'_μ and $W_{\mu 4}$. It is noteworthy that in the limit $v_{1,2} = 0$, the elements $M_{24}'^2$ and $M_{34}'^2$ vanish, and there is no mixing between W_4 and Z_μ, Z'_μ .

In the second step, three remain neutral gauge bosons gain masses via seesaw mechanism:

$$M_Z^2 = \frac{g^2}{4} [M_{22}'^2 - (M^{off})^T (M_{2 \times 2}'^2)^{-1} M^{off}], \tag{93}$$

where

$$M^{off} = \begin{pmatrix} M_{23}'^2 \\ M_{24}'^2 \end{pmatrix}, \quad M_{2 \times 2}'^2 = \begin{pmatrix} M_{33}'^2 & M_{34}'^2 \\ M_{34}'^2 & M_{44}'^2 \end{pmatrix}. \tag{94}$$

Combining (93) and (94) yields

$$\begin{aligned}
M_Z^2 &= \frac{g^2}{4} \left(M_{22}'^2 + \frac{(M_{24}'^2)^2 M_{33}'^2 - 2M_{23}'^2 M_{24}'^2 M_{34}'^2 + (M_{23}'^2)^2 M_{44}'^2}{(M_{34}'^2)^2 - M_{33}'^2 M_{44}'^2} \right) \\
&= \frac{g^2 (u^2 + u'^2 + v^2 + v'^2)}{2c_W^2} - \frac{g^2}{2c_W^2} \Delta_{M_z^2},
\end{aligned}$$

where

$$\begin{aligned}
\Delta_{M_z^2} &= \frac{4\Delta_z^2 (4c_W^4 x_3 - x_0 x_1 + x_4) + x_1 [x_2 x_1 - 4\Delta_z^2 x_0]}{x_2 (x_4 + 4c_W^4 x_3) - 4\Delta_z^2 x_0^2} \\
&= \frac{4\Delta_z^2 (4c_W^4 x_3 - 2x_0 x_1 + x_4) + x_1^2 x_2}{x_2 (x_4 + 4c_W^4 x_3) - 4\Delta_z^2 x_0^2},
\end{aligned} \tag{95}$$

with

$$\begin{aligned}
x_1 &= (1 - 2c_W^2)(u^2 + u'^2) + v^2 + v'^2, \\
x_2 &= 2\Lambda_1(2\lambda_1 + \Lambda_1) + 2\Lambda_2(2\lambda_2 + \Lambda_2) + \omega^2 + u^2 + u'^2, \\
x_3 &= 4\Lambda_1^2 + 4\Lambda_2^2 + \omega^2 + u^2 + u'^2, \quad x_4 = (1 - 4c^2)(u^2 + u'^2) + v^2 + v'^2, \\
\Delta_z &= \Lambda_1 v_1 + \Lambda_2 v_2.
\end{aligned}$$

The ρ parameter in our model is given by

$$\rho = \frac{M_W^2}{M_Z^2 \cos^2 \theta_W} = 1 + \delta_{\text{tree}}, \quad (96)$$

where

$$\delta_{\text{tree}} = \frac{\delta_{WZ}}{M_Z^2}, \quad \delta_{WZ} = \frac{g^2}{2c_W^2} (\Delta_{M_Z^2} - \Delta_{M_W^2}). \quad (97)$$

Using approximations (74) and (75), we have

$$\begin{aligned} \Delta_{M_Z^2} - \Delta_{M_W^2} &\simeq 8(\Lambda_1 v_1 + \Lambda_2 v_2) \left\{ -\frac{\Lambda_2 v_2}{2\Lambda_1^2 + 2\Lambda_2^2 + \omega^2} \right. \\ &+ \left. \frac{(4\Lambda_1^2 + 4\Lambda_2^2 + \omega^2)(4c_W^2 - 1)c_W^2(\Lambda_1 v_1 + \Lambda_2 v_2)}{2(4c_W^2 - 1)[(2\Lambda_1^2 + 2\Lambda_2^2 + \omega^2)(4\Lambda_1^2 + 4\Lambda_2^2 + \omega^2)c_W^4 - (4c_W^2 + 1)^2(\Lambda_1 v_1 + \Lambda_2 v_2)^2]} \right\}. \end{aligned} \quad (98)$$

We assume relations (57) and $v_2 \equiv v_s$, $\omega = \Lambda_2 \equiv \Lambda_s$; then

$$\begin{aligned} \Delta_{M_Z^2} - \Delta_{M_W^2} &\simeq 8(k^2 + 1)\Lambda_s v_s \left[-\frac{v_s}{(2k^2 + 3)\Lambda_s} + \frac{(k^2 + 1)(4k^2 + 5)c_W^2 \Lambda_s v_s}{2[(8k^4 + 22k^2 + 15)c_W^4 \Lambda_s^2 - (k^2 + 1)^2(4c_W^2 + 1)^2 v_s^2]} \right] \\ &\simeq -\frac{8(k^2 + 1)v_s^2}{2k^2 + 3} + \frac{8(k^2 + 1)^2(4k^2 + 5)c_W^2 v_s^2}{2(2k^2 + 3)(4k^2 + 5)c_W^4} \\ &\simeq -\frac{8(k^2 + 1)v_s^2}{2k^2 + 3} + \frac{8(k^2 + 1)^2 v_s^2}{2(2k^2 + 3)c_W^2} \\ &= \frac{8(k^2 + 1)v_s^2}{2k^2 + 3} \left(\frac{k^2 + 1}{2c_W^2} - 1 \right). \end{aligned} \quad (99)$$

$$\Delta_{M_Z^2} - \Delta_{M_W^2} \simeq \frac{8(k^2 + 1)v_s^2}{2k^2 + 3} \left(\frac{k^2 + 1}{2c_W^2} - 1 \right). \quad (100)$$

From (97) and (100), we have

$$\delta_{\text{tree}} = \frac{g^2}{2c_W^2} \frac{1}{M_Z^2} \frac{8(k^2 + 1)v_s^2}{2k^2 + 3} \left(\frac{k^2 + 1}{2c_W^2} - 1 \right). \quad (101)$$

The experimental value of the ρ parameter and M_W are given in Ref. [4]:

$$\begin{aligned} \rho &= 1.0004_{-0.0004}^{+0.0003} \quad (\delta_{\text{tree}} = 0.0004_{-0.0004}^{+0.0003}), \\ s_W^2 &= 0.23116 \pm 0.00012, \\ M_W &= 80.358 \pm 0.015 \text{ GeV}. \end{aligned} \quad (102)$$

Hence,

$$0 \leq \delta_{\text{tree}} \leq 0.0007. \quad (103)$$

From (102) and (103), we can deduce the relations between v, g and k . Indeed,

$$v = \pm \frac{c_W^2 \sqrt{\delta_{\text{tree}}} \sqrt{2k^2 + 3M_Z}}{g \sqrt{2k^2 + 2} \sqrt{k^2 + 1 - 2c_W^2}}$$

The Fig. 1 gives the relation between v_s and g, k with $g = 0.5$ and $k \in (0.9, 1.1)$, for $|v_s| \in (0, 8)$ GeV. Conditions (74) and (75) are then satisfied. The Fig. 2 gives the relation between g and $\delta_{\text{tree}}, v_s$




FIG. 1: The relation between v_s and g, k with $g = 0.5$ and $k \in (0.9, 1.1)$

with $k = 1$ and $\delta_{\text{tree}} \in (0, 0.0007)$, $v_s \in (0, 8.0)$ GeV, for $|g| \in (0, 2)$ GeV. Conditions (74) and (75) are then satisfied. The Fig. 3 gives the relation between k and g, v_s with $\delta_{\text{tree}} = 0.0005$ and




FIG. 2: The relation between g and $\delta_{\text{tree}}, v_s$ with $k = 1$ and $\delta_{\text{tree}} \in (0, 0.0007)$, $v_s \in (0, 8.0)$ GeV

$g \in (0.4, 0.6)$, $v_s \in (0, 8.0)$ GeV, for $k \in (1, 3)$ GeV (k is a real number, Fig. 3a) or $k = ik_1, k_1 \in (-1.2, -1.05)$ GeV (k is a pure complex number, Fig. 3b). Conditions (74) and (75) are then

satisfied. From Fig. 3 we see that many values of k that are different from close to unity still can fit the recent experimental data [4]. This means that the difference of $\langle s_1 \rangle$ and $\langle s_1 \rangle$ as mentioned in this work is necessary.




FIG. 3: The relation between k and g, v_s provided $\delta_{\text{tree}} = 0.0005$ and $g \in (0.4, 0.6)$, $v_s \in (0, 8.0)$ GeV

Diagonalizing the mass matrix $M_{2 \times 2}'^2$, we obtain two new physical gauge bosons

$$\begin{aligned} Z_\mu'' &= \cos \phi Z_\mu' + \sin \phi W_{\mu 4}, \\ W_{\mu 4}' &= -\sin \phi Z_\mu' + \cos \phi W_{\mu 4}. \end{aligned} \quad (104)$$

The mixing angle ϕ is given by

$$\tan \phi = \frac{4\sqrt{\alpha_0}c_W(\Lambda_1 v_1 + \Lambda_2 v_2)x_0}{-4c_W^4\alpha_0 x_3 + c_W^2 x_2 - \alpha_0 x_4 + \sqrt{F}}, \quad (105)$$

where

$$F = \left(4c_W^4\alpha_0 x_3 - c_W^2 x_2 + \alpha_0 x_4\right)^2 + 16\alpha_0 c_W^2 (\Lambda_1 v_1 + \Lambda_2 v_2)^2 x_0^2.$$

If $\lambda_{1,2}^2, v_{1,2}^2, u^2, u'^2, v^2, v'^2 \ll \omega^2 \sim \Lambda_s^2 \sim \Lambda_\sigma^2$ then

$$\sqrt{F} \simeq c_W^2 [2\Lambda_1^2 + 2\Lambda_2^2 + \omega^2 - 4\alpha c_W^2 (4\Lambda_1^2 + 4\Lambda_2^2 + \omega^2)],$$

and we can evaluate

$$\tan \phi \simeq -\frac{2\sqrt{\alpha_0}(\Lambda_1 v_1 + \Lambda_2 v_2)x_0}{c_W [2(8\alpha_0 c_W^2 - 1)(\Lambda_1^2 + \Lambda_2^2) + (4\alpha_0 c_W^2 - 1)\omega^2]} \sim \frac{v_i}{\Lambda_i} \quad (i = 1, 2). \quad (106)$$

The physical mass eigenvalues are defined by

$$M_{Z_\mu'', W_{\mu 4}'}^2 = \frac{g^2}{4c_W^2} \left\{ 4\alpha_0 c_W^4 x_3 + c_W^2 x_2 + \alpha_0 x_4 \pm \sqrt{F} \right\}. \quad (107)$$

In the limit $\lambda_{1,2}, v_{1,2} \rightarrow 0$, the mixing angle ϕ tends to zero, and $M_{Z''_\mu, W'_{\mu 4}}^2$ in (107) reduces to

$$\begin{aligned} M_{Z''_\mu}^2 &= \frac{g^2}{2c_W^2} \left[c_{2W}^2(u^2 + u'^2) + v^2 + v'^2 + 4c_W^4(4\Lambda_1^2 + 4\Lambda_2^2 + \omega^2) \right] \alpha_0, \\ M_{W'_{\mu 4}}^2 &= \frac{g^2}{2} (u^2 + u'^2 + \omega^2 + 2\Lambda_1^2 + 2\Lambda_2^2). \end{aligned} \quad (108)$$

From (69) and (108), the $W'_{\mu 4}$ and $W_{\mu 5}$ components have the same mass, and hence, in this approximation we should identify the linear combination

$$\sqrt{2}X_\mu^0 = W'_{\mu 4} - iW_{\mu 5} \quad (109)$$

as a physical neutral non-Hermitian gauge boson. The subscript "0" indicates the neutrality of the gauge boson X_μ . We note that the identification in (109) can only be acceptable in the limit $\lambda_{1,2}, v_{1,2} \rightarrow 0$. In general, it is not true because of the difference in masses of $W'_{\mu 4}$ and $W_{\mu 5}$ as in (68) and (107).

Expressions (79) and (106) show that, with the limits (74) and (75), the mixings between the charged gauge bosons $W - Y$ and the neutral ones $Z' - W_4$ are of the same order because they are proportional to v_i/Λ_i ($i=1,2$). In addition, from (108)

$$M_{Z''_\mu}^2 \simeq g^2(4\Lambda_1^2 + 4\Lambda_2^2 + \omega^2)$$

is somewhat bigger than

$$M_{W'_{\mu 4}}^2 \simeq \frac{g^2}{2} (\omega^2 + 2\Lambda_1^2 + 2\Lambda_2^2)$$

(or $M_{X_\mu^0}^2$), and

$$|M_Y^2 - M_{X_\mu^0}^2| = \frac{g^2}{2} (u^2 + u'^2 - v^2 - v'^2)$$

is slightly smaller than

$$M_W^2 = \frac{g^2}{2} (u^2 + u'^2 + v^2 + v'^2).$$

In that limit, the masses of X_μ^0 and Y degenerate.

VI. CONCLUSIONS

We have studied new features of the 3-3-1 model with a neutral fermion based on the S_3 flavor symmetry in which the anti-sextet responsible for neutrino mass and mixing lies in the 2

representation under S_3 and the number of Higgs multiplets required is reduced. If the S_3 symmetry is violated as a perturbation by the difference in components of the anti-sextet, S_3 is equivalently broken into identity, the corresponding neutrino mass mixing matrix acquires the most general form. This way of the symmetry breaking helps us reduce the content in the Higgs sector: only one anti-sextet instead of three multiplets (two anti-sextets and one triplet) as in our previous work. By assuming that the VEVs of the anti-sextet differ from each other and regarding the difference between these VEVs as a small perturbation, we can make the model fit the latest data on neutrino oscillations. Our results show that the neutrino masses are naturally small and a deviation from the tri-bimaximal neutrino mixing form can be realized. The Higgs potential of the model and minimization conditions are also considered.

Acknowledgments

This work was supported in part by the National Foundation for Science and Technology Development of Vietnam (NAFOSTED) under grant number 103.01-2011.63.

-
- [1] Super-Kamiokande collaboration, Y. Fukuda *et al.* (1998), Phys. Rev. Lett. 81, 1562, arXiv: 9807003 [hep-ex]; Super-Kamiokande collaboration, S. Fukuda *et al.* (2001), Phys. Rev. Lett. 86, 5651, arXiv: 0103032 [hep-ex]; Super-Kamiokande collaboration, S. Fukuda *et al.* (2001), Phys. Rev. Lett. 86, 5656, arXiv: 0103033 [hep-ex]; Super-Kamiokande collaboration, S. Fukuda *et al.* (2002), Phys. Lett. B 539, 179, arXiv: 0205075 [hep-ex].
 - [2] Super-Kamiokande collaboration, Y. Ashie *et al.* (2004), Phys. Rev. Lett. 93, 101801, arXiv: 0404034 [hep-ex]; Super-Kamiokande collaboration, Y. Ashie *et al.* (2005), Phys. Rev. D 71, 112005, arXiv: 0501064 [hep-ex].
 - [3] K. Nakamura *et al.* [Particle Data Group], J. Phys. G **37**, 075021 (2010).
 - [4] J. Beringer, *et al.* (2012), *Review of Particle Physics* (Particle Data Group), Phys. Rev. D. 86, 010001; T. Schwetz, M. Tortola, and J. Valle (2011), New J. Phys. 13, 109401, arXiv: 131108.1376 [hep-ph]; K. Abe *et al.* (2011) [T2K Collaboration], Phys. Rev. Lett. 107, 041801; P. Adamson *et al.* (2011) [MINOS Collaboration], Phys. Rev. Lett. 107, 181802, arXiv: 1108.0015 [hep-ex]; Fogli, G.L. *et al.* (2011), Phys. Rev. D 84, 053007, arXiv: 1106.6028 [hep-ph]
 - [5] A. Zee, Phys. Lett. B 93, 389 (1980); P. H. Frampton and S. L. Glashow, Phys. Lett. B 461, 95 (1999); A. S. Joshipura, S. D. Rindani, Phys. Lett. B 464 239 (1999); Y. Koide Nucl. Phys. B (Proc. Suppl.)

- 111, 294 (2002), ; K. Hasegawa, C. S. Lim, K. Ogure , Phys. Rev. D **68**, 053006 (2003), arXiv: 0303252 [hep-ph].
- [6] K. S. Babu, Phys. Lett. B **203**,132 (1988).
- [7] F. Pisano and V. Pleitez, Phys. Rev. D **46**, 410 (1992); P. H. Frampton, Phys. Rev. Lett. **69**, 2889 (1992); R. Foot, O. F. Hernandez, F. Pisano and V. Pleitez, Phys. Rev. D **47**, 4158 (1993).
- [8] A. Doff and F. Pisano, Mod. Phys. Lett. A **15**, 1471 (2000); P. V. Dong and H. N. Long, Int. J. Mod. Phys. A **21** 6677, (2006).
- [9] W. A. Ponce, Y. Giraldo and L. A. Sanchez, Phys. Rev. D **67**, 075001 (2003); P. V. Dong, H. N. Long, D. T. Nhung and D. V. Soa, Phys. Rev. D **73**, 035004 (2006); P. V. Dong and H. N. Long, Adv. High Energy Phys. **2008**, 739492 (2008), [arXiv:0804.3239(hep-ph)].
- [10] M. Singer, J. W. F. Valle and J. Schechter, Phys. Rev. D **22**, 738 (1980); J. C. Montero, F. Pisano and V. Pleitez, Phys. Rev. D **47**, 2918 (1993); R. Foot, H. N. Long and Tuan A. Tran, Phys. Rev. D **50**, 34(R) (1994), [arXiv: hep-ph/9402243]; H. N. Long, Phys. Rev. D **53**, 437 (1996); H. N. Long, Phys. Rev. D **54**, 4691 (1996).
- [11] P. V. Dong, L. T. Hue, H. N. Long and D. V. Soa, Phys. Rev. D **81**, 053004 (2010).
- [12] P. V. Dong, H. N. Long, D. V. Soa, and V. V. Vien, Eur. Phys. J. C **71**, 1544 (2011).
- [13] P. V. Dong, H. N. Long, C. H. Nam, V. V. Vien (2012), Phys. Rev. D **85**, 053001, arXiv: 1111.6360 [hep-ph].
- [14] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530**, 167 (2002); Z. Z. Xing, Phys. Lett. B **533**, 85 (2002); X. G. He and A. Zee, Phys. Lett. B **560**, 87 (2003); Phys. Rev. D **68**, 037302 (2003).
- [15] E. Ma and G. Rajasekaran, Phys. Rev. D **64**, 113012 (2001); K. S. Babu, E. Ma and J. W. F. Valle, Phys. Lett. B **552**, 207 (2003); G. Altarelli and F. Feruglio, Nucl. Phys. B **720**, 64 (2005); E. Ma, Phys. Rev. D **73**, 057304 (2006); X. G. He, Y. Y. Keum and R. R. Volkas, JHEP **0604**, 039 (2006); S. Morisi, M. Picariello, and E. Torrente-Lujan, Phys. Rev. D **75**, 075015 (2007); C. S. Lam, Phys. Lett. B **656**, 193 (2007); F. Bazzocchi, S. Kaneko and S. Morisi, JHEP **0803**, 063 (2008); A. Blum, C. Hagedorn, and M. Lindner, Phys. Rev. D **77**, 076004 (2008); F. Bazzochi, M. Frigerio, and S. Morisi, Phys. Rev. D **78**, 116018 (2008); G. Altarelli, F. Feruglio and C. Hagedorn, JHEP **0803**, 052 (2008); M. Hirsch, S. Morisi and J. W. F. Valle, Phys. Rev. D **78**, 093007 (2008); E. Ma, Phys. Lett. B **671**, 366 (2009); G. Altarelli and D. Meloni, J. Phys. G **36**, 085005 (2009); Y. Lin, Nucl. Phys. B **813**, 91 (2009); Y. H. Ahn and C. S. Chen, Phys. Rev. D **81**, 105013 (2010); J. Barry and W. Rodejohanny, Phys. Rev. D **81**, 093002 (2010); 119901(E) (2010); G. J. Ding and D. Meloni, Nucl. Phys. B **855**, 21 (2012).
- [16] R. N. Mohapatra, M. K. Parida, G. Rajasekaran, Phys. Rev. D **69**, 053007 (2004); C. Hagedorn, M. Lindner, and R. N. Mohapatra, JHEP **0606**, 042 (2006); E. Ma, Phys. Lett. B **632**, 352 (2006); H. Zhang, Phys. Lett. B **655**, 132 (2007); Y. Koide, JHEP **0708**, 086 (2007); H. Ishimori, Y. Shimizu, and M. Tanimoto, Prog. Theor. Phys. **121**, 769 (2009); F. Bazzocchi, L. Merlo, and S. Morisi, Nucl. Phys. B **816**, 204 (2009); F. Bazzocchi and S. Morisi, Phys. Rev. D **80** 096005 (2009); G. Altarelli and F. Feruglio, Rev. Mod. Phys. **82**, 2701 (2010); G.J. Ding, Nucl. Phys. B **827**, 82 (2010); Y. H. Ahn, S.

- K. Kang, C. S. Kim, and T. P. Nguyen, Phys. Rev. D **82**, 093005 (2010); Y. Daikoku and H. Okada, arXiv:1008.0914 [hep-ph]; H. Ishimori, Y. Shimizu, M. Tanimoto, and A. Watanabe, Phys. Rev. D **83** 033004 (2011); H. Ishimori and M. Tanimoto, Prog. Theor. Phys. **125**, 653 (2011); R. Z. Yang and H. Zhang, Phys. Lett. B **700** 316, (2011); S. Morisi and E. Peinado, Phys. Lett. B **701**, 451 (2011); S. Morisi, K. M. Patel, and E. Peinado, Phys. Rev. D **84**, 053002 (2011).
- [17] L. Wolfenstein, Phys. Rev. D **18**, 958 (1978); S. Pakvasa and H. Sugawara, Phys. Lett. B **73**, 61 (1978); **82**, 105 (1979); E. Durman and H. S. Tsao, Phys. Rev. D **20**, 1207 (1979); Y. Yamanaka, H. Sugawara, and S. Pakvasa, Phys. Rev. D **25**, 1895 (1982); K. Kang, J. E. Kim, and P. Ko, Z. Phys. C **72**, 671 (1996); H. Fritzsch and Z. Z. Xing, Phys. Lett. B **372**, 265 (1996); K. Kang, S. K. Kang, J. E. Kim, and P. Ko, Mod. Phys. Lett. A **12**, 1175 (1997); M. Fukugita, M. Tanimoto, and T. Yanagida, Phys. Rev. D **57**, 4429 (1998); H. Fritzsch and Z. Z. Xing, Phys. Lett. B **440**, 313 (1998); Y. Koide, Phys. Rev. D **60**, 077301 (1999); H. Fritzsch and Z. Z. Xing, Phys. Rev. D **61**, 073016 (2000); M. Tanimoto, Phys. Lett. B **483**, 417 (2000); G. C. Branco and J. I. Silva-Marcos, Phys. Lett. B **526**, 104 (2002); M. Fujii, K. Hamaguchi and T. Yanagida, Phys. Rev. D **65**, 115012 (2002); J. Kubo, A. Mondragon, M. Mondragon and E. Rodriguez-Jauregui, Prog. Theor. Phys. **109**, 795 (2003); **114**, 287(E) (2005); P. F. Harrison and W. G. Scott, Phys. Lett. B **557**, 76 (2003); S.-L. Chen, M. Frigerio, and E. Ma, Phys. Rev. D **70**, 073008 (2004); **70**, 079905(E) (2004); H. Fritzsch and Z. Z. Xing, Phys. Lett. B **598**, 237 (2004); F. Caravaglios and S. Morisi, arXiv:hep-ph/0503234; W. Grimus and L. Lavoura, JHEP **0508**, 013 (2005); R. N. Mohapatra, S. Nasri and H. B. Yu, Phys. Lett. B **639**, 318 (2006); R. Jora, S. Nasri and J. Schechter, Int. J. Mod. Phys. A **21**, 5875 (2006); J. E. Kim and J.-C. Park, JHEP **0605**, 017 (2006); Y. Koide, Eur. Phys. J. C **50**, 809 (2007); A. Mondragon, M. Mondragon, and E. Peinado, Phys. Rev. D **76**, 076003 (2007); AIP Conf. Proc. **1026**, 164 (2008); M. Picariello, Int. J. Mod. Phys. A **23**, 4435 (2008); C. Y. Chen and L. Wolfenstein, Phys. Rev. D **77**, 093009 (2008); R. Jora, J. Schechter and M. Naeem Shahid, Phys. Rev. D **80**, 093007 (2009); **82**, 079902(E) (2010); D. A. Dicus, S. F. Ge and W. W. Repko, Phys. Rev. D **82**, 033005 (2010); Z. Z. Xing, D. Yang and S. Zhou, Phys. Lett. B **690**, 304 (2010); R. Jora, J. Schechter and M. N. Shahid, Phys. Rev. D **82**, 053006 (2010); S. Dev, S. Gupta and R. R. Gautam, Phys. Lett. B **702**, 28 (2011); D. Meloni, S. Morisi and E. Peinado, J. Phys. G **38**, 015003 (2011); G. Bhattacharyya, P. Leser and H. Pas, Phys. Rev. D **83**, 011701(R) (2011); T. Kaneko and H. Sugawara, Phys. Lett. B **697**, 329 (2011); S. Zhou, Phys. Lett. B **704**, 291 (2011).
- [18] A. E. C. Hernandez, R. Martinez and J. Nisperuza, *S₃ flavour symmetry breaking scheme for understanding the quark mass and mixing pattern in SU(3)_C ⊗ SU(3)_L ⊗ U(1)_X models*, e-Print: arXiv:1401.0937 [hep-ph].
- [19] V. V. Vien and H. N. Long, *Int. J. Mod. Phys. A* **28**, 1350159 (2013), [arXiv:1312.5034 (hep-ph)].
- [20] E. Ma, arXiv:hep-ph/0409075; H. Ishimori *et. al.*, Prog. Theor. Phys. Suppl. **183**, 1 (2010).
- [21] V. D. Barger, S. Pakvasa, T. J. Weiler and K. Whisnant, Bimaximal mixing of three neutrinos, Phys. Lett. B **437**, 107 (1998), arXiv: 9806387 [hep-ph]; I. Stancu and D. V. Ahluwalia, Phys. Lett. B **460**, 431 (1999), arXiv: 9903408 [hep-ph]; H. Georgi and S. L. Glashow, Phys. Rev. D **61**, 097301 (2000),

- arXiv: 9808293 [hep-ph]; N. Li and B.-Q. Ma, Phys. Lett. B600, 248 (2004), arXiv: 0408235 [hep-ph].
- [22] P. V. Dong, V. T. N. Huyen, H. N. Long, and H. V. Thuy, Gauge Boson Mixing in the 3-3-1 Models with Discrete Symmetries, *Advances in High Energy Physics*, **2012**, 715038 (2012).
- [23] P. V. Dong and H. N. Long, Eur. Phys. J. C 42, 325 (2005).